

University of Louisville

ThinkIR: The University of Louisville's Institutional Repository

Electronic Theses and Dissertations

12-2003

Functional proteomic analysis of the myocardial PKCepsilon subproteome PKCepsilon-AKT-eNOS signaling modules and during.

Jun Zhang
University of Louisville

Follow this and additional works at: <https://ir.library.louisville.edu/etd>

Recommended Citation

Zhang, Jun, "Functional proteomic analysis of the myocardial PKCepsilon subproteome PKCepsilon-AKT-eNOS signaling modules and during." (2003). *Electronic Theses and Dissertations*. Paper 1635.
<https://doi.org/10.18297/etd/1635>

This Doctoral Dissertation is brought to you for free and open access by ThinkIR: The University of Louisville's Institutional Repository. It has been accepted for inclusion in Electronic Theses and Dissertations by an authorized administrator of ThinkIR: The University of Louisville's Institutional Repository. This title appears here courtesy of the author, who has retained all other copyrights. For more information, please contact thinkir@louisville.edu.

**FUNCTIONAL PROTEOMIC ANALYSIS OF THE MYOCARDIAL PKC ϵ
SUBPROTEOME AND PKC ϵ -AKT-eNOS SIGNALING MODULES
DURING CARDIOPROTECTION**

By

Jun Zhang
M.S., Chinese Academy Of Sciences, 1994

A Dissertation
Submitted to the Faculty of the
Graduate School of the University of Louisville
In Partial Fulfillment of the Requirements
For the Degree of

Doctor of Philosophy

Department of Physiology & Biophysics
University of Louisville
Louisville, Kentucky

December, 2003

**Functional Proteomic Analysis Of The Myocardial PKC ϵ Subproteome
And PKC ϵ -Akt-eNOS Signaling Modules
During Cardioprotection**

By

Jun Zhang
M.S., Chinese Academy of Sciences, 1994

A Dissertation Approved on

Sept. 18th, 2003
Date

By the following Dissertation Committee:

Dissertation Director

DEDICATION

*For my parents
and my dear husband Lizhi*

ACKNOWLEDGMENTS

I would like to express my deep appreciation to my mentor Dr. Peipei Ping for her extraordinary guidance, invaluable support and precious friendship. Her example of hard work, dedication and courage have been a great inspiration.

I also wish to express my truly gratitude to the members of my graduate advisory committee, Dr. Aruni Bhatnagar, Dr. Stanley E. D' Souza, Dr. Irving G. Joshua, Dr. John C. Passmore, Dr. William B. Wead for providing their generous support and excellent advise throughout my Ph.D. studies. Additional, many thanks to the rest of the faculty and administration of the Department of Physiology and Biophysics who have been of tremendous help during my graduation.

I also like to extend my appreciation to other members (current and former) of the PKC laboratory for all their support and friendship: Dr. Thomas M Vondriska, Dr. Christopher Baines, Mr. Ernest M. Cardwell, Dr. Guangwu Wang, Mr. Nobel Zong, Dr. Xinan Cao, Dr. Yuting Zheng, Dr. Changxu Song, Dr. Xin Qiao, Ms. Dawn Pantaleon. In Particular, I sincerely thank Dr. Thomas M Vondriska and Dr. Christopher Baines for their excellent instruction.

Finally, I would like to express my special thanks to my parents for their unconditional love, encouragement, and great support. Also, I can not thank enough my dear husband Lizhi Fu and my son for their boundless love, encouragement and tremendous support.

Thank you all !

ABSTRACT

FUNCTIONAL PROTEOMIC ANALYSIS OF THE MYOCARDIAL PKC ϵ SUBPROTEOME AND PKC ϵ -AKT-eNOS SIGNALING MODULES DURING CARDIOPROTECTION

Jun Zhang

September, 2003

Ischemic heart disease is the leading cause of mortality and morbidity in Western societies. Thus, understanding the molecular mechanisms to reduce myocardial ischemia and limit infarction size are of great importance. Ample evidence has shown that protein kinase C epsilon (PKC ϵ) play an essential role in the genesis of cardioprotection. In particular, our laboratory has shown that activation of PKC ϵ in the heart is sufficient to significantly reduce myocardial infarction due to coronary artery occlusion. However, the molecular mechanism responsible for PKC ϵ -induced cardioprotection remain unclear. Recently, functional proteomic analysis have demonstrated that PKC ϵ forms signaling complexes with various of proteins, including structural proteins, signaling molecules, stress-activated proteins, transcriptional/translational factors, metabolism-related proteins, and PKC-binding domain containing proteins, and the coordination of these molecules contributes to cardioprotection. Despite this information which obtained from whole heart lysates, little is known regarding the cellular mechanisms that regulate the assembly of these protein complexes and the specific manner in which these molecules interact with each. Therefore, this dissertation is focused on the characterization of PKC ϵ subproteome and signaling modules within, as well as their regulation during cardioprotection. PKC ϵ -Akt-eNOS signaling module is chosen as my research paradigm because the individual molecule PKC ϵ , Akt and eNOS has

been implicated in the prevention of cell death. In addition, both eNOS and Akt were found to exist in the PKC ϵ complex in our initial studies, suggesting that functional coupling among these molecules may be a heretofore unrecognized protective signaling mechanism. Our data demonstrate that the cardiac PKC ϵ signaling subproteome is comprised of various sizes of protein complexes, and the formation of these complexes and modules is regulated by the subcellular localization and extracellular stimulus. During cardioprotection, activation of PKC ϵ enhances the PKC ϵ -Akt-eNOS module formation, and thus contributes to the regulation of nitric oxide production and hence the manifestation of the cardioprotective phenotype. In conclusion, the present research successfully characterizes the myocardial PKC ϵ subproteome and PKC ϵ -Akt-eNOS signaling modules, which serves as an important step towards our complete understanding of the signaling mechanisms underlying PKC ϵ -mediated cardioprotection and ultimately aids in the development of pharmacological agents to protect the heart.

TABLE OF CONTENTS

	PAGE
ACKNOWLEDGEMENTS	iv
ABSTRACT	v
LIST OF FIGURES	ix
 CHAPTER I. GENERAL INTRODUCTION AND OVERVIEW	
1. Proteome, subproteome, and module	1-4
2. Ischemic injury and cardioprotection	5
3. Protein kinase C epsilon (PKC ϵ)	6-9
4. Akt/PKB	10-13
5. Endothelial nitric oxide synthase (eNOS)	14-16
6. PKC ϵ -Akt-eNOS signaling modules	17-18
 CHAPTER II: METHODS AND MATERIALS	33-38
 CHAPTER III: Functional proteomic mapping of the myocardial PKCϵ signaling	
subproteome in the mouse heart: a subcellular compartment study.	
Abstract	39-40
Introduction	41-42
Results	43-49
Discussion	50-53
 CHAPTER VI: Functional proteomic analysis of myocardial PKCϵ-Akt-eNOS signaling	
modules during cardioprotection.	
Abstract	75-76
Introduction	77-78
Results	79-82
Discussion	83-87
 REFERENCES	102-115

CURRICULUN VITAE 116-122
-------------------------	---------------

LIST OF FIGURES

FIGURES	PAGE
1. Functional proteomic platform for isolation and characterization of multiprotein complexes	19
2. Molecular structure of Protein kinase C epsilon (PKC ϵ)	21
3. Model of activation of PKC ϵ	23
4. Activation of PKC ϵ reduces myocardial infarction sizes	25
5. Molecular structures and activation model of Akt isoforms.....	27
6. Molecular structures of endothelial nitric oxide synthase (eNOS).	29
7. Architecture of the cardioprotective PKC ϵ signaling system in cardiac myocytes	31
8. Flow chart for isolation and visualization of PKC ϵ signaling multiprotein complexes (experimental strategy for Chapter III)	54
9. Separation protein mixtures from cytosolic and membrane fractions by two dimensional electrophoresis	56
10. Immunoblotting analysis of known marker proteins in subcellular fractions	58
11. Characterization of multiprotein PKC ϵ signaling complexes-A	60
12. Characterization of multiprotein PKC ϵ signaling complexes-B	62
13. Characterization of multiprotein PKC ϵ signaling complexes-C	64
14. Demonstration of multiprotein PKC ϵ signaling complexes by liquid chromatography	66
15. Determination of the native size of multiprotein PKC ϵ complexes	68
16. Identification of the molecular components of the multiple PKC ϵ complexes: examples of eNOS and p170 PI 3-kinase	70

17.	PKC ϵ directly interact with Akt and eNOS <i>in vitro</i>	88
18.	Activation of both PKC ϵ and Akt enhanced PKC ϵ -Akt-eNOS modules formation <i>in vitro</i>	90
19.	Formation of PKC ϵ -Akt-eNOS signaling modules <i>in vivo</i>	92
20.	PKC ϵ phosphorylated and enhanced activation of Akt and eNOS <i>in vitro</i>	94
21.	PKC ϵ directly phosphorylated Akt at its Ser473 site <i>in vitro</i>	96
22.	Effect of PKC ϵ on the phosphorylation and activation of Akt <i>in vivo</i>	98
23.	Effect of PKC ϵ on the phosphorylation of eNOS <i>in vivo</i>	100

CHAPTER I

General Introduction And Overview

Proteome, subproteome, and module

Upon completion of the human genome project, we have entered the era of proteomics. Proteomics is the study of the proteome, which was first defined in 1995 as all proteins expressed by the genome of an organism or tissue (175). Despite the power of genomic technologies, and the information derived from a sequenced genome, it is clear that proteins are ultimate functional units within cells, and the coordination of them determines the biological phenotype. Therefore, mapping the human proteome will bridge the gap between genomics and systems physiology, and provide tremendous information for diagnostic and therapeutic applications.

The goal of proteomics is to globally map protein expression profiles and protein-protein interaction profiles within the whole organism (133, 134). However, the generation of proteome map faces great challenges. First, one gene does not encode a single protein. Indeed, the number of proteins in the entire human body is expected to be many times higher than that of genes (81, 96). Second, concomitant co- and post-translational modification events, such as phosphorylation, glycosylation, sulphonation, palmitoylation, hydroxylation, acetylation and myristolation et al., which play critical roles in cellular function, can further increase the complexity of protein products from a single reading frame. For example, it is estimated that 30% of all cellular proteins contain covalently bound phosphate (94), and phosphorylation events control the majority of the signal transduction pathways in eukaryotic cells. Third, in contrast to the static genome, the proteome is a dynamic entity that is constantly being modulated by biological and environmental stimuli (74, 86). Fourth, the architecture of a proteome is both cell and organ specific. Finally, although significant advances have been made in the development of practical

proteomic techniques, more high-throughput platforms need to be evolved to enhance the power of current approaches (95). Taken together, mapping the entire proteome is a long-term project due to its enormity and complexity.

Instead of characterizing every protein within the cell, the goal of *functional* proteomic analysis is to analyze the spatial and temporal properties of the specific sets of proteins, called subproteomes in a cell, tissue, or organism, which are related to the particular biological functions (73, 170, 173). Compared with the traditional proteomic analysis, functional proteomic studies are more feasible and advantageous. Proteins are not separate islands that perform their functions entirely by themselves. Instead, mounting evidence supports the concept that a protein may assembly of multi-protein complexes with other proteins to serve as a means to accomplish complicated tasks of cellular function (3). Characterization of the specific subproteome provides information about the components of the protein complex, and its dynamic changes, including protein expression, modification, and subcellular localization, in response to different conditions. So far, several studies (6, 22, 66, 71, 87, 98, 99, 101, 113, 132, 155) have undertaken the characterization of different subproteomes, including those defined by the MAPK cascade, the NMDA receptor complexes, nucleoporin complexes, and the PDGF receptor signaling system. Within the subproteome, two or more proteins are associated to form a basic functional unit of signal transduction, called a module, which conducts different tasks at various subcellular compartments. The formation of many distinct modules contributes to the establishment specific biological phenotype (170, 172).

A technical platform has been developed in our lab for comprehensive analysis of functional subproteomes (Figure 1) (135, 170, 171). In order to characterize proteins that associate with a protein of interest, isolation and purification of protein complexes is the first step. After subcellular fractionation which is an effective way to reduce the complexity and enrich the low-abundance proteins, immunoprecipitation and recombinant protein-based affinity pull down assays are commonly used to isolate protein complexes. Protein complexes are then separated by one- or two-dimensional electrophoresis, or liquid chromatography, followed by mass spectrometry or Western blotting to identify components within each complex. Several techniques

aid the further study of identification proteins, such as confocal microscopy to confirm the subcellular localization, biochemical *in vitro* assay to test the protein-protein interaction, and protein assays to detect the protein expression profiles within the subproteome. Finally, determination of the functional role of individual protein within the subproteome can link proteomic data to a physiological phenotype. This step can be conducted by using cell culture or transgenic animal models, or treatment of activators or inhibitors to test the biological significance of the protein change in response to the changed environmental conditions.

Similarly, the cardiac proteome is proposed to compose of multiple integrated functional subproteomes, each of which is organized to facilitate a unique function. Moreover, specific modules are formed within subproteomes in response to different stimuli (145, 160, 172). Instead of attempting to simultaneously examine all proteins in cardiac cells, we focus our research on characterization myocardial subproteomes that are critically involved in the development of ischemic injury and cardioprotection, especially the subproteomes of protein kinases associated with cardioprotection (136, 142, 172). Three aspects are important to characterize a subproteome: the identification of the proteins that compose the subproteome are classified according to their biological functions; the molecular architecture of the subproteome, the manner in which the subproteome is altered in different subcellular locations; and the dynamic modulation of the subproteome that underlies the development of a specific phenotype. Delineation of multiple subproteomes based on their respective biological functions and contributions to phenotypes will eventually lead to full characterization of cardiac proteome.

The goal of this dissertation was to contribute to this understanding of cell signaling in two fundamental regards. First, as detailed in Chapter III, we wanted to examine the role of subcellular-dependent regulation of signal transduction by testing the basic hypothesis that the cell recruits different proteins to regions of the cell to perform specific tasks, and that multiprotein complexes are involved in this phenomenon. Second, as reported in Chapter IV, we targeted investigation of a specific signaling module within the cardiomyocyte to test a separate but equally central hypothesis, namely, that the cell organizes multiple molecules into signal transduction units (or modules) that are mobilized to accomplish signaling events and to regulate

phenotype. In this regard, the present dissertation advances our understanding of cell signaling in general on two critical levels that are interrelated to the common theme of PKC ϵ -mediated cardioprotection.

Ischemic injury and cardioprotection

Ischemic heart disease is the leading cause of mortality and morbidity in Western societies (25). Thus, mechanisms to reduce myocardial ischemia and to prevent myocytes death are of great importance to improve patient survival. Due to the limited ability of the heart to regenerate cardiomyocytes, myocardial cell death is the most deleterious event in the heart. In the past, numerous attempts had been undertaken to prevent myocardial cell death and limit the infarct size after a coronary artery occlusion. An endogenous protective defense mechanism termed ischemic preconditioning (PC) was discovered in 1986 (128), and exerts a very powerful anti-infarct effect. Ischemic PC is a phenomenon whereby brief ischemic episodes activate an innate protective system in the heart to render the heart powerfully resistance to subsequent prolonged lethal ischemic insults. Over the past 16 years, a tremendous evidence has demonstrated that this protective phenomenon exists in virtually all mammalian models(11, 23). It is hoped that elucidation of the intracellular signaling mechanisms underlying the myocardial preconditioning may lead to delivery of effective therapeutic agents capable of reducing the myocardial infarction. Importantly, cardioprotection can also be engendered by the administration of pharmacological agents such as nitric oxide (NO) donors *in lieu of* ischemia (140, 179, 181). NO donor-induced cardioprotection may have potential clinical applications.

Mounting evidence has demonstrated that protein kinase C epsilon plays a critical role in the genesis of cardioprotection (49, 75, 117, 143, 186). In particular, cardiac-specific activation of PKC ϵ affords an infarct-sparing effect similar in efficacy to that induced by ischemic and pharmacological PC. However, the molecular and cellular mechanisms underling PKC ϵ -induced cardioprotection are poorly understood.

Protein kinase C epsilon (PKC ϵ)

The protein kinase C (PKC) family of serine-threonine kinases is composed of 12 different isoforms that are divided into three subgroups based on their regulation. The classical or conventional PKCs (cPKCs) include the α , β I, β II, γ isoforms, and are regulated by Ca^{2+} , diacylglycerol (DAG), and phosphatidylserine (PS). The novel PKCs (nPKCs), which include the ϵ , δ , η , and θ isoforms, are regulated by DAG and PS; but are insensitive to Ca^{2+} . The atypical PKCs (aPKCs) include the ζ , ι , and λ isoforms and are regulated by PS alone. The μ isoform of PKC (also known as PKD), which is sensitive to both DAG and PS, has been placed into a separate class because of its unique structural characteristics (129) (Figure 2).

Molecular structure of PKC ϵ : Protein kinase C ϵ (PKC ϵ) was first cloned in 1987 and defined as a novel PKC isoform. Akin to other PKCs isoforms, PKC ϵ consists of two primary functional domains: an amino-terminal regulatory domain and a carboxyl-terminal catalytic domain (129) (Figure 2). Within the regulatory domain, there are two C1 regions and one C2 region. The C1 region contains C1A and C1B which form the DAG/phorbol esters binding site via the cysteine-rich Zn^{2+} finger motif. An autoinhibitory pseudosubstrate sequence precedes the C1 domain which can autoinhibit PKC ϵ to form an inactive closed conformation by occupying the catalytic substrate-binding cavity. Unlike cPKCs, the C2 region of PKC ϵ lacks the specific aspartate residues which contribute to the Ca^{2+} binding. Therefore, activation of PKC ϵ is not dependent upon the concentration of Ca^{2+} ions. The catalytic domain has an ATP-binding site (C3) and a substrate-binding site (C4), both of which are essential for kinase activity. The hinge region between C2 and C3 domain can be easily cleaved to release the carboxyl terminal constitutively active enzyme (kinase domain) when PKC ϵ is membrane-bound (Figure 3).

Regulation of PKC ϵ : There are at least three conserved phosphorylation sites within PKC ϵ that are necessary for the maturation of PKC ϵ and its ability to respond to second messengers. The first, rate-limiting step is the phosphorylation of Thr566 in the activation loop (kinase core). Phosphoinositide-dependent kinase-1 (PDK-1) may be involved in this phosphorylation event that leads to the exposure of C-terminus of PKC to the following two phosphorylation modifications via intracellular autophosphorylation or catalysis by other kinases (33, 161). These two phosphorylation sites are Thr710 at the turn motif (164), and Ser729 in hydrophobic site in C-terminus. Phosphorylation of these three sites results in conformational changes of PKC ϵ that not only lock PKC ϵ into a more thermally stable, protease- and phosphatase-resistant stage, but also free the substrate-binding cavity for pseudosubstrate to form the closed conformation (Figure 3).

Mature but inactive PKC ϵ localizes in the cytosolic fraction where it is maintained in the autoinhibited closed conformation. PKC ϵ can be activated by several different second messengers, including, DAG, PIP₃ (phosphatidylinositol 3,4,5-triphosphate), and fatty acids that are produced by physiological stimuli such as PDGF (platelet-derived growth factor) and bradykinin (72, 127). After stimulation, PKC ϵ binds to membrane-associated DAG via its C1 domain. This stimulates the translocation of PKC ϵ from cytosolic fractions to membrane fractions where C2 region can interact with membrane phosphatidyl serine (PS) (129). This tethering of both C1 and C2 regions to membrane lipids provide the energy to release the pseudosubstrate motif from the catalytic domain and open the enzyme's conformation, resulting in maximal PKC ϵ activation.

Subcellular distribution of PKC ϵ : The subcellular distribution behavior of PKC ϵ depends on (i) which second messenger is bound to C1 domain and (ii) which intracellular compartment the PKC ϵ -specific binding partner, called RACK2 (also known as β' -COP), is localized. This latter molecule binds the C2 (-like) domain of PKC ϵ and modulates its activity. Therefore, both the C1 domain and C2 domain of PKC ϵ function as subcellular localization signals (112). The spatial-

and temporal- regulation of PKC ϵ subcellular localization may provide the explanation for its substrate specificity in the signaling transduction.

Function of PKC ϵ in non-cardiac cells: PKC ϵ is expressed in many tissues and cells, with particular abundance in neuronal, hormonal, and immune cells. PKC ϵ has been reported to play essential roles in a variety of cellular signaling processes (2), including proliferation, differentiation, gene expression, muscle contraction, mechanical force adaptation, metabolism, transport, exocytosis, endocytosis, and inflammation. PKC ϵ deficient mice show macrophage dysfunction due to decreased activation of the transcription factor NF- κ B, resulting in increased mortality after bacterial infection, and indicating the important protective role of PKC ϵ in the immune system (32). In the neuronal system, PKC ϵ induces neurite outgrowth during neuronal differentiation through its interaction with actin filaments (106). PKC ϵ has also been implicated in the regulation of nociceptor function that participates in the perception of pain (46). Moreover, PKC ϵ could promote cell survival and prevent apoptosis through suppression of the excitotoxic glutamate receptor in neuronal PC12 cells (167), and through dysregulation of the mitochondrial caspase pathway in lung cancer cells (46).

PKC ϵ and cardioprotection: In cardiac cells, multiple investigators have found that activation and translocation of PKC ϵ serves as a central signaling event in the development of cardiac protection (75, 117, 140-142). In the past several years, studies from our laboratory (140-142) have shown that cardiac-specific overexpression of PKC ϵ in transgenic mice hearts can mimic protection against myocardial infarction by ischemic preconditioning (Figure 4). Studies based on the cardio-specific transgenic expression of the PKC ϵ -selective translocation inhibitor (epsilonV1) or activator (ψ epsilon RACK) peptides from other groups also clearly show the importance of PKC ϵ in the cardiac protection against ischemic damage as well as during normal postnatal myocardial development (50). Furthermore, moderate ethanol consumption exerts beneficial cardioprotective effects via mechanisms that involve PKC ϵ activation (188).

Although PKC ϵ has an established role in the genesis of cardioprotection, the molecular and cellular mechanisms responsible for PKC ϵ -mediated cardiac protection remain unclear.

Mounting evidence suggests that PKC-dependent protection involves a battery of kinases (tyrosine kinases, Akt, PI3-K, mitogen-activated protein kinase (MAPKs)) (145, 146, 166), stress-activated proteins (heat shock proteins (HSPs), nitric oxide synthases (NOSs), ion channels, and pro-survival transcription factors (NF- κ B, HIF-1 α). Recently, using functional proteomic approaches, our laboratory has demonstrated that PKC ϵ forms signaling complexes with multiple proteins and that the coordinated action of these molecules confers protection against ischemic injury (12, 20, 21, 130). In other words, there is a cardiac signaling network centered around PKC ϵ that is critical for cardioprotection.

AKT / Protein Kinase B

Akt, also called protein kinase B (PKB), is a serine/threonine kinase, which was first discovered in 1991 by two independent research groups. Akt was initially characterized as the human cellular homologue of the viral oncogene *v-akt* from the transforming retrovirus AKT8 (19, 38, 100). To date, three major isoforms of Akt/PKB have been identified, encoded by three distinct gene loci, termed Akt1/ α , Akt2/ β , and Akt3/ γ . All three isoforms share a high degree of homology at the amino acid level (42, 103). Akt2 and Akt3 are approximate 82% identical with the Akt1 isoform. Akt3 lacks 23 amino acids at the C-terminal compared with others (110) (Figure 5). All three isoforms are ubiquitously expressed in mammals, although levels of expression vary among tissues. Akt1 is the predominant isoform in most tissues, such as brain, thymus, heart and lung (19, 38, 100). Akt2 is primarily expressed in the insulin-responsive tissues, such as skeletal muscle, heart, liver and kidney (7). Expression of Akt3 is more restricted, appearing high in the brain and testes, and lower in heart, spleen, lung and skeletal muscle. All tissues contain at least one form of Akt.

Molecular structure of Akt/PKB: Examination of the amino acid sequence of Akt/PKB revealed that all three isoforms contain a highly conserved structure. The N-terminal 100 amino acids possess a pleckstrin homology (PH) domain that mediates lipid-protein or/and protein-protein interactions. A short glycine-rich region links the PH domain to the catalytic domain. The central kinase catalytic domain has a conserved phosphorylation site Thr308 (Akt2 Thr309, Akt3 Thr302) within the activation loop. In C-terminal regulatory domain there is another conserved phosphorylation site Ser473 (Akt2 Ser474, Akt3 Ser472) (Figure 5).

Regulation of Akt/PKB: In the inactive form of Akt, the PH domain serves as an inhibitory structure that sterically blocks the catalytic site (similar to the autoinhibition of PKC ϵ). Activation of Akt is initiated by the binding of PIP₃ to the PH domain of this enzyme. This action not only

releases autoinhibition through conformational changes that unmask the active site, but also recruits Akt to the plasma membrane where it can be further phosphorylated and fully activated. So far, phosphorylation of Thr308 in the kinase activation loop and Ser473 in the C-terminal hydrophobic region are the two most important posttranslational modifications. Mutation of Thr308 to nonphosphorylatable alanine abolishes Akt 's activation, indicating that phosphorylation of Th308 is essential for activation of Akt. In contrast, mutation of Ser473 only partially inhibits Akt activation, suggesting that Ser473 phosphorylation facilitates maximal Akt activation but is not a necessary step in Akt-dependent signaling (5, 138, 183).

Ser124 and Thr450, which are constitutively phosphorylated, were also predicted to be prerequisite for Akt responsiveness (18), although, mutation of these residues to alanine has no effect on the activity of Akt (7, 165). In addition, a recent report finds that Src family tyrosine kinase dependent phosphorylation of two tyrosine residues (Y315, and Y326) may contribute to the activation of Akt (35).

The cellular activation of Akt/PKB is dependent upon the interaction with membrane-bound second messenger PIP_3 . The generation of PIP_3 is catalyzed by PI-3K via the phosphorylation of phosphoinositides (PtdIns). Although PIP_3 does not activate Akt/PKB directly, it recruits Akt to the plasma membrane and alters the conformation of Akt to facilitate the subsequent phosphorylation modifications. The upstream kinase that phosphorylates Thr308 is a 67kDa ubiquitously expressed kinase called PDK-1 (5, 162, 163). The role of PDK-1 to activate Akt has been supported by studies from embryonic stem cells deficient in PDK-1 (177), biochemical inhibition of PDK-1 (13), and by studies employing anti-sense oligonucleotides against PDK-1 (59). Akt/PKB becomes a substrate for PDK-1 once its PH domain engages PIP_3 on the membrane. Interestingly, as indicated above, PDK-1 is also known to phosphorylate and activate PKC, though the mechanisms contributing to the specificity of PDK-1 for PKC, Akt, or other substrates, remain poorly defined.

The upstream kinase for Ser473 phosphorylation is unknown, and is tentatively named "PDK-2". Several candidate activities have been proposed, including $PKC\zeta$ (34, 190), an

unnamed lipid-raft-associated kinase (84), or simply an autophosphorylation processes following the PDK-1 activity (1).

There are two postulated models for Akt activation. The first mechanism in response to PI3K induction and PIP₃ production, Akt moves from the cytosol to the plasma membrane, where Ser473 phosphorylation occurs through autophosphorylation or by PDK2. This region forms a docking site for PDK-1 to phosphorylate Thr308 and hence activate Akt (8). The second mechanism is that cytosolic Akt and PDK-1 are co-localized, but Akt is inactive due to constraints imposed by its PH domain. Growth factors stimulate PI3K, which draw both PDK-1 and Akt to the plasma membrane. The PH domain of Akt binds to PIP₃, unmasking the activation loop and allowing PDK-1 to phosphorylate Thr308. Subsequent elevation of Akt activity promotes autophosphorylation of Ser473 or by a third-party enzyme (165). Once Akt is activated, it targets specific proteins in the cytoplasm, plasma membrane and and nucleus (Figure 5).

The consensus motif for Akt phosphorylation is R-x-R-x-x-S/T-F/L (x represents any amino acid) (5). So far, several potential Akt substrates have been found, such as glycogen synthase kinase-3 (GSK3), I κ B- α kinase α (IKK α), Raf, and Bad. Recently, Akt was found to regulate the level of nitric oxide (NO) production through direct phosphorylation and activation of the endothelial isoform of nitric oxide synthase (eNOS) (44, 63).

Subcellular distribution of Akt/PKB: The subcellular localization of Akt is tightly regulated. Two minutes after stimulation by IGF-1, Akt1 is translocated to the plasma membrane in a PH domain-dependent manner. Prolonged stimulation induces Akt translocated to the nucleus by an unknown mechanism (8), which is PH-domain-independent.

Function of Akt/PKB: Ample evidence shows that Akt is a key player in numerous physiological and pathophysiological processes, such as gene transcription, protein synthesis, glycogen synthesis, transcription control, cell growth, cell survival and angiogenesis (26, 28, 44, 62, 63, 67, 69, 121,182). In cardiomyocytes, Akt is activated by gp130 stimulation and has both a hypertrophic and an antiapoptotic effect (62, 131). Bell and colleagues reported that administration of bradykinin, as an adjunct to reperfusion, limits infarct size in the mouse heart via induction of increased activity of PI3kinase, Akt and eNOS (62). In myocardial gene transfer

studies, a constitutively active mutant of Akt reduces infarct size and apoptosis after ischemia-reperfusion injury (62). Using DNA microarrays, Cook et al (40) reported the cardiac gene expression profiles in mice with cardiac-specific overexpression of Akt. Studies from these Akt mice also indicate that Akt exerts a dual function *in vivo* on the myocardium by increasing both cardiomyocyte size and contractility (39). They found that chronic Akt activation resulted in the differential regulation of at least 40 genes in the heart that may contribute to the effects of Akt on cardiomyocyte survival, metabolism, and growth. Interestingly, studies from Zhou's group find that moderate alcohol consumption induces sustained cardiac protection is through a PKC ϵ - and Akt-dependent signaling pathway (188).

Endothelial nitric oxide synthase (eNOS)

Nitric oxide (NO) is a potent messenger molecule that participates in many physiological processes such as vascular relaxation, neuronal communication, immunological modulation, and host defense (4). NO is produced from oxidizing L-arginine to generate L-citrulline which is catalyzed by a family of NO synthases (NOS). There are three distinct isoforms of NOS, referred to as endothelial NOS (eNOS, NOS-3), neuronal NOS (nNOS, NOS-1), and inducible NOS (iNOS, NOS-2). eNOS and nNOS are Ca^{2+} /calmodulin-dependent enzymes producing NO in short bursts and low concentrations (nM), whereas iNOS produces NO in high concentrations (μM) and is Ca^{2+} -independent (9).

Molecular structure of eNOS: eNOS was first identified and isolated from bovine aortic endothelial cells (9, 10) in 1991. Since then, it has been found in many cells, such as platelets (148), smooth muscle cells (156), cardiac myocytes (14), bone cells, (4) and neurons (27, 45). According to the amino acid sequence, the primary structure of eNOS has two functional domains: reductase domain and oxygenase domain. The C-terminal reductase domain (10) contains NADPH, FAD, FMN, and calmodulin (CaM) binding sites and the N-terminal oxygenase domain has binding sites for tetrahydrobiopterin (BH₄), heme, and L-arginine (14, 27). In particular, the BH₄ domain is reported to be critical for eNOS activity (116, 169), as it is involved in dimer formation and electron transfer, as well as maintenance of the active site's structural integrity (Figure 6).

The quaternary structure of eNOS includes its dimerization and association with other proteins. The eNOS enzyme is the only fully functional in a dimeric form. Dimerization of eNOS starts with the binding of heme (108), and the binding of BH₄, to stabilize the dimer (102, 116, 150). The reductase domain transfers electrons from NADPH via the FAD, FMN and CaM co-factors to the heme in the oxygenase domain. The oxygenase domain catalyzes L-arginine to L-

citrulline with BH₄ binding to O₂. CaM binding increases the rate of electron transfer, by which increased intracellular Ca²⁺ concentration is required for this protein-protein interaction. Hence, the Ca²⁺/CaM-dependence is one of the features of this particular NOS isoform (Figure 6).

Regulation of eNOS: The regulation of eNOS occurs at three levels: gene transcription, co/post-translational modification, and allosteric regulation of the enzyme itself.

Transcriptional/translational control alters eNOS protein expression levels via regulation of the eNOS promoter and altered stability of eNOS mRNA. There are numerous transcription factor binding sites on the eNOS promoter, including activator protein (AP)-1, nuclear factor (NF)-1, nuclear factor- κ B (NF- κ B). Multiple stimuli can affect the basal expression of eNOS. For example, hypoxia leads to the activation of AP-1 and induction of eNOS transcription (53, 89). eNOS levels in endothelial cells also can be efficiently regulated by changes in eNOS mRNA stability. Hypoxia and cytokines, such as TNF, can downregulate eNOS mRNA levels by reducing the half-life of eNOS mRNA from 48 h to as little as 3 h (122, 125, 185, 187).

There are several phosphorylation modifications within the eNOS protein. Ser1177 (human) in the reductase domain is the most important one. It is stimulated by the application of fluid shear stress (44, 64), estrogen (111), VEGF (107), insulin (58), or bradykinin (16) and results in increased NO production. The kinases involved in this process vary with the stimuli and include Akt (44, 51), PKA (123), and CaMKII (120). In addition, Thr495 within the CaM-binding is constitutively phosphorylated in endothelium cells, possibly via PKC, and is associated with decreased eNOS activity (43, 85, 120, 123). Moreover, other phosphorylation sites, such as Ser114 within the oxygenase domain, are thought to be modified by ERK1/2 (56) and are related to the dimerization of eNOS. Ser633 and some tyrosine residues can also be phosphorylated, but the functional significance of these post-translational modifications remains to be determined (44, 56, 57).

Lastly, the molecular chaperone Hsp90 serves as an allosteric modulator (65) for eNOS activity. Activation of vascular endothelial growth factor, histamine or fluid shear stress in human endothelial cell increases the interaction between eNOS and Hsp90, and increases eNOS activity by approximate 3-fold.

Subcellular localization of eNOS: eNOS is largely membrane-associated as a result of N-terminal myristoylation and palmitoylation modifications (30, 152, 157). When eNOS is located in the caveolae, flask-shaped invaginations of the plasma membrane that occupy up to 30% of cell surface, it is bound to caveolin. Caveolin is a caveolar resident coat protein that inhibits eNOS activity by interfering with Ca^{2+} /CaM binding and electron transfer to the heme subunits (9). In endothelial cells, eNOS binds to caveolin-1, whereas in cardiac myocytes, eNOS is associated with caveolin-3. An increase in intracellular Ca^{2+} causes the formation of Ca^{2+} /CaM complexes, which displaces caveolin resulting in the dissociation of eNOS. At this point, eNOS is now activated and remains so until intracellular Ca^{2+} levels decrease and the Ca^{2+} /CaM complex can no longer compete with caveolin. The specific intracellular compartments to which eNOS translocates remain unclear, but many include the nuclear membrane, the endoplasmic reticulum, or trans-golgi system (83).

Function of eNOS: The protective effects of NO generated by eNOS are varied. It regulates vascular tone by relaxation of vascular smooth muscle cells (vasodilation) (76). It inhibits leukocyte and platelet adhesion and aggregation to endothelium (100) and inhibits vascular smooth muscle cell proliferation (156).

eNOS is the major form of NOS in the heart and it is expressed in endothelial cells and cardiac myocytes (14, 105). The protective role of eNOS in the cardiovascular system has been widely studied. Myocardial overexpression of eNOS in mice greatly stimulates cardiac NO formation and protects the heart against ischemia/reperfusion injury (29). Moreover, gene transfer of eNOS to the heart reduces ischemia/reperfusion injury by inhibiting NF- κ B activation, adhesion molecule expression, and the early infiltration of leukocytes (92). In contrast, eNOS knockout mice exhibit increased left ventricular contractility in response to β -adrenergic agonists compared with wide type mice, demonstrating a negative inotropic effect of eNOS-derived NO (15, 70, 79). eNOS is also reported to be involved in the development of ischemic preconditioning. Xuan's group report that in conscious rabbit hearts, ischemic PC elicits an immediate activation of Ca^{2+} -dependent NOS (most likely eNOS), indicating that eNOS serves as a trigger on day 1 to initiate the development of late PC 24 hours later (181).

PKC ϵ -Akt-eNOS signaling modules

As detailed above, PKC ϵ plays a critical role in the genesis of cardioprotection against myocardial ischemic injury (49, 140-143, 147). Specifically, recent studies in our laboratory showed that PKC ϵ forms multimeric signaling complexes with at least 93 proteins, including structural proteins, stress-activated proteins, proteins associated with transcription and translation, proteins involved in cellular metabolism, and various kinases and phosphatases (52, 143). Furthermore, the assembly of these multiprotein PKC ϵ complexes is dynamically modulated in the PKC ϵ -transgenic mouse hearts, which serves as important evidence for the role of these complexes in regulating the cardioprotective phenotype.

Importantly, among the 93 known protein partners, Akt and eNOS are of great interest as they are well established to independently promote cell survival and prevent cell death in response to various forms of stress (16, 42, 60, 63, 104, 124, 182). Taken in the context of the foregoing introduction regarding the essential roles of these three molecules in the cardiovascular system, the finding that PKC ϵ , Akt and eNOS may be signaling partners in the myocardium was a tantalizing piece of evidence suggesting functional coupling of these molecules. As a result, we chose to focus our studies to define the specific role of subcellular dependent regulation of the PKC ϵ subproteome, and in particular, of PKC ϵ -Akt-eNOS signaling modules within this subproteome.

Accordingly, my research has focused on the identification of PKC ϵ subproteome, and the signaling modules within. We hypothesized that PKC ϵ form distinct signaling modules at different subcellular compartments with different protein components in the mouse heart, and moreover, that the assembly and activation of signaling modules conduct different tasks which are essential to the genesis of cardioprotection (Figure 7). In the first part of my research, I concentrated on the identification of PKC ϵ subproteome in different subcellular compartments,

and the regulation of these protein complexes during cardioprotection. In the second part of my study, I focused on the characterization of PKC ϵ -Akt-eNOS signaling modules, which were chosen as an example to functionally validate a signaling module in cardioprotection.

Previous studies in non-cardiac cells have provided evidence for physical and functional interaction between PKC, Akt and eNOS. Akt can directly interact and phosphorylate eNOS on Ser1177 (63). Doornbos' (48) and Mao's (119) groups reported that Akt can be co-immunoprecipitated with PKC ζ or PKC ι , and that the PH domain of Akt is implicated as the binding site for PKC ζ to Akt1 (110) and Akt3 (88) in COS-1 cells. Li and colleagues also find that PKC α overexpression stimulates Akt activity and suppresses apoptosis induced by interleukin 3 withdrawals in myeloid progenitor cells (114). In contrast, Wen (176) reported that in A549 and HEK293 cells, the phosphatidylinositol 3-kinase/Akt signaling pathway is negatively regulated by PKC. Similarly, PMA-induced apoptosis is accompanied by an inhibition of Akt activity (149). Moreover, PKC activation inhibited eNOS activity by attenuating eNOS phosphorylation on Ser1177 and increasing phosphorylation of Thr495 in cardiovascular endothelial cells (123, 124).

Despite this information, nothing is known regarding the formation and regulation of PKC ϵ -Akt-eNOS signaling modules in myocytes, especially during cardioprotection. Therefore, comprehensive investigations need to be implemented to draw a clearer picture of the temporal and spatial relationship among PKC ϵ , Akt and eNOS in the regulation of NO production and the genesis of cardioprotection. Successful characterization of myocardial PKC ϵ -Akt-eNOS signaling modules would serve as a good model for the future research addressing the signaling mechanisms underlying PKC ϵ -mediated cardioprotection, and of critical importance in the development of pharmacological agents for the treatment of myocardial ischemia and infarction in patients. This dissertation provided new information about the molecular architecture of PKC ϵ -Akt-eNOS signaling module, the subcellular localization of these modules, and the dynamic modulation of these modules during the development of the cardioprotective phenotype.

Figure 1. Functional proteomic platform for isolation and characterization of multiprotein complexes. This is a hypothesis-driven research strategy to characterize a cellular subproteome responsible for the genesis of a phenotype. Started with subcellular fractionation which is an effective way to reduce the complexity and enrich the low-abundance proteins, immunoprecipitation and recombinant protein-based affinity pull down assays are commonly used to isolate and purify protein complexes. Protein complexes are then separated by one- or two-dimensional electrophoresis, or liquid chromatography, followed by mass spectrometry or Western blotting to identify components within each complex. Then, biochemical analyses and phenotype assessment are conducted to determine the functional role of individual protein within the subproteome and link proteomic data to a physiological phenotype. We use this strategy to characterize the PKC ϵ -dependent signaling mechanisms underlying cardioprotection.

Functional Proteomic Technology Platform




Figure 2. Molecular structure of protein kinase C epsilon (PKC ϵ). Alike other PKCs, PKC ϵ has regulatory domain and kinase domain. Within the regulatory domain, an autoinhibitory pseudosubstrate sequence is ahead of C1 motif (dotted red). 2nd messenger DAG and phorbol esters bind to C1 domain; whereas, acidic lipids (PS) binds to C2 domain. Kinase core has conserved three phosphorylation sites. N is the amino terminus. C is the carboxyl terminus.




Figure 3. Model of activation of PKC ϵ . Newly synthesized immature PKC ϵ need three conserved phosphorylation modifications to become maturation which is phosphorylated by PDK-1 followed by autophosphorylation. Mature but inactive PKC ϵ localizes in cytosolic fraction. In response to various stimulus, the generation of DAG and PS recruits PKC ϵ to the membrane fraction by engaging the C1 and C2 domains on the membrane. This membrane interaction provides the energy to release the pseudosubstrate motif from the substrate-binding cavity, resulting open confirmation and maximal PKC ϵ activation.




Figure 4. Activation of PKC ϵ reduces myocardial infarction sizes. PKC ϵ transgenic mice was generated using α -MHC cardiac-specific promoter. The DNA constructs used in the generation of constitutively active PKC ϵ mutant transgenic mice has a point mutation at the pseudosubstrate region (A to E at aa159). This mutation would release pseudosubstrate domain from catalytic site, thus placing PKC ϵ in its open, and therefore active conformation. When these mice were subjected to myocardial ischemia, the PKC ϵ transgenic mice show significantly decreased myocardial infarction size when compared with non-transgenic mice, indicating that active PKC ϵ protect the heart.





Figure 5. Molecular structures and activation model of Akt/PKB isoforms. Upper panel, Akt is also a serine/threonine kinases with three major isoforms. Each isoform consists of an N-terminal PH domain, a kinase domain and a C-terminal regulatory domain. N-terminal PH domain can mediate lipid-protein or protein-protein interactions. Central kinase domain has a conserved phosphorylation site Thr308 which is necessary for Akt activation. The hydrophobic C-terminal regulatory domain containing a second regulatory phosphorylation site Ser473, which is required for maximal Akt activation. Lower panel, inactive Akt mainly resides in the cytosol, and the PH domain is serving as an autoinhibitory motif to block kinase domain. (like PS for PKC ϵ). The activation of Akt requires the binding of PIP₃ to PH domain to unmask the kinase domain, leading to the translocation to membrane fraction; Akt activation also requires the phosphorylation modifications of Thr308 and Ser473, which is mediated by kinase called PDK1 and unknown kinases called PDK-2.





Figure 6. Molecular structure of endothelial nitric oxide synthase (eNOS). eNOS is Ca^{2+} -CaM-dependent nitric oxide synthases, and has two functional domains. C-terminal reductase domain and N-terminal oxygenase domain. The myristoylation (Myr) and palmitoylation (Palm) sites on eNOS are shown (important features for eNOS membrane localization), and the autoinhibitory loop within the FMN regions of eNOS is also shown. eNOS can catalyze arginine to produce nitric oxide. And its activity is highly regulated by phosphorylation of Ser1177, resulting in an increased activity and increased NO production.





Figure 7 Architecture of the cardioprotective PKC ϵ signaling system in cardiac myocytes. The hypothetical structure of the PKC ϵ signaling subproteome involves various signaling modules located at multiple subcellular compartments, and their integrity carries out signal transduction tasks required for cardioprotection.



PKCε

PKCε Signaling modules

PKCε subproteome

Phenotype

CHAPTER II

METHODS AND MATERIALS

All procedures were performed in accordance with the Animal Research Committee guidelines at UCLA, which conform with the *Guide for the Care and Use of Laboratory Animals*, published by the National Institutes of Health.

Antibodies and Chemicals

Anti- β '-adaplin, BMX, caveolin-3, connexin-43, COX-2, desmin, ERK1, HIF-1 α , GM130, Lap2, MEK1, eNOS, HSP60, iNOS, NuMA, p38 MAPK, p85 PI-3K, p170 PI-3K, PKC ϵ , Pyk2, Ras, RACK1, and villin antibodies were purchased from Pharmingen (San Diego, CA). Anti-AKAP79, Akt, ANT1, c-Fos, c-Jun, c-Myc, CREB, Elk, HO-1 HSP27, HSP70, GRP75, hnRNP α 2/b1, JNK2, lamin-C, Lck, NF- κ B (p65), PDK1, Src, STAT1 and 3, Tropinin I and T, vimentin, vinculin, Akt1/2, Akt1, Akt2 and Akt3 antibodies were purchased from Santa Cruz Biotechnology (Santa Cruz, CA). Anti- α B-crystallin and TCP- α (RACK2) antibodies were purchased from Stressgen (Vancouver, BC). Anti-phospho-Akt of Ser308 and Ser 473, anti-phospho-eNOS (Ser1177) antibodies and Akt kinase assay kit were from Cell Signaling Technology Inc. (Beverly, MA). Anti- α -actin and α -actinin antibodies from Sigma-Aldrich (St. Louis, MO). Anti-Na⁺/K⁺ ATPase, histone H4, PP2A, PP1 antibodies were purchased from Upstate Biotechnology (Lake Placid, NY), anti-prohibitin-1 antibody from Research Diagnostics (Flanders, NJ), and anti-VDAC1 from calbiochem (San Diego, CA). Anti-ribonucleoprotein (RNP) antibody was from NeoMarkers (Fremont, CA), anti-glyceraldehydes-3-PDH (GAPDH) antibody was from Biodesign (Saco, Maine), and anti-creatine kinase was from abcam (Cambridge, UK). The aldose reductase antibody was a generous gift from Dr. Aruni Bhatnagar (University of Louisville). The hnRNPH1

antibody and hnRNP K antibody were generous gifts from Dr. Doug Black (Howard Hughes Medical Institutes, CA), and Dr. Pradip Raychaudhuri (University of Illinois at Chicago), respectively. The Oracle antibody was custom made by BioSource (Camarillo, CA). Cross-linker EZ-Link Sulfo-NHS-LC-Biotin and streptavidin:HRP were obtained from Pierce (Rockford, IL). L-arginine, Bistris, tricine, sucrose, 6-aminocapric acid, and other chemicals were purchased from Sigma-Aldrich. Recombinant active Akt, inactive Akt, PH domain of Akt, PH domain deleted Akt were purchased from Upstate Biotechnology (Lake Placid, NY). Recombinant PKC ϵ and eNOS were bought from Biomol (Plymouth meeting, PA) and Calbiochem (La Jolla, CA), respectively.

PKC ϵ Transgenic Mice

The cardiac-targeted PKC ϵ transgenic mice used in these studies have been previously described (142, 145). This transgenic line exhibits moderate increases in PKC ϵ activity, is inherently cardiac protected, and is free from hypertrophy. Transgenic mice and their non-transgenic littermates were used at 9-12 weeks.

Subcellular fractionation

All fractionation procedures were performed at 4°C. Myocardial cytosolic, nuclear, mitochondrial, membrane fractions were obtained by differential centrifugation as previously described (12). Briefly, mouse hearts were homogenized in buffer containing 250mM sucrose, 10mM Tris-HCl (pH7.4), 1mM EDTA, 1mM Na₃VO₄, and a cocktail of protease inhibitors (Roche, Indianapolis, IN). The homogenate was then centrifuged at 1000 x g for 10 min to pellet the nuclear fraction. The supernatant was centrifuged at 10,000 x g for 30 min to yield the mitochondrial fraction in the pellet (12). The resultant supernatant was subsequently centrifuged at 100,000 x g for 1 hr to yield the cytosolic fraction (supernatant) and membrane fraction (pellet). The different membrane fractions were then re-suspended in homogenization buffer with 0.5% (v/v) NP-40. Caveolae were isolated by differential density centrifugation, as described by Smart *et al* (159).

Immunoprecipitation

PKC ϵ monoclonal antibodies were covalently cross-linked to protein-A/G beads (Santa Cruz) using a commercially available kit (Pierce). Subcellular lysates (1000 μ g) were incubated with the immobilized PKC ϵ antibody overnight at 4°C and washed three times with buffer containing 150 mM NaCl, 20 mM Tris-HCl (pH 7.4), 10 mM EDTA, 1 % (v/v) NP-40, 1 mM Na₃VO₄, and a protease inhibitor cocktail (Roche). After the final wash, the beads were re-suspended in Laemmli buffer, boiled for 10 min, and then the supernatants were subjected to SDS-PAGE and immunoblotting. If immunoprecipitation was followed by two dimensional (2D) electrophoretic analysis, after the final wash, the beads were washed with 10% sucrose to remove salt, and the protein complexes then dissolved into thiourea buffer (Urea 7M, thiourea 2M, CHAPS 2% (w/v), DTT 65mM, Zwittergent 1% (v/v), Ampholytes (pH3-10) 0.8% (w/v), and bromphenol blue 0.01% (w/v) (Genomic Solutions, Ann Arbor, MI).

Immunoblotting

Samples were denatured in Laemmli buffer, resolved by one-dimensional SDS-PAGE, and transferred onto nitrocellulose membranes. After blocking with 5% (w/v) nonfat milk powder, the membranes were immunoblotted using the ECL detection system (Amersham, Piscataway, NJ).

Two-Dimensional Electrophoresis

Subcellular extracts were loaded onto immobilized pH gradient (IPG) strips (Amersham) and subjected to isoelectric focusing in the first dimension as described (142). The IPG strips were re-equilibrated and subjected to SDS-PAGE (Genomic Solutions) for the second dimension. Proteins were then visualized by silver staining.

Protein Biotinylation

Following PKC ϵ immunoprecipitation, protein complexes were eluted with low pH elution buffer (Pierce) and then dialyzed against PBS overnight at 4°C. To biotinylate the proteins, the dialyzed sample was incubated with EZ-Link Sulfo-NHS-LC-Biotin (Pierce) at 4°C for 4 hours. The

samples were then subjected to SDS-PAGE, transferred to nitrocellulose membranes, and visualized using streptavidin-HRP (Pierce) and the ECL system (Amersham).

Gel Filtration Chromatography

Gel filtration chromatography was carried out as described (36) with a few modifications. Subcellular fractionations which were prepared from 15 hearts in Buffer A containing: 20mM HEPES, pH 7.9, 1.5mM MgCl₂, 150mM NaCl, 0.2mM EDTA, 0.5% (v/v) NP-40, and a cocktail of protease inhibitors, or total homogenates from two mouse hearts were dialyzed overnight at 4°C in Buffer A without NP-40, and clarified prior to chromatography. These samples were then loaded onto a pre-calibrated Sephacryl S-400 column (XK 26/70, 26 mm diameter, 70 cm length, Amersham) using Buffer A without NP-40 as running buffer. Each 1ml fraction was collected. Proteins were then precipitated with cold acetone, and analyzed by either SDS-PAGE or native gel electrophoresis. Thyroglobulin (669kDa), Ferritin (440kDa), catalase (232kDa) and aldolase (158kDa) (Pharmacia) were used as molecular standards.

Native Gel Electrophoresis

Native gel electrophoresis was carried out as described (153, 154) with minor modifications. In particular, deoxycholic acid was used in place of Coomassie blue. Gel filtration fractions were precipitated with cold acetone and then solubilized in buffer containing 50mM 6-aminocaproic acid, 200mM Bis-Tris, pH 7.0, 20% (v/v) glycerol, and 0.5% (v/v) NP-40 for 1 hour on ice. Then samples were mixed with 10x native sample buffer containing 100mM Bis-Tris, pH 7.0, 1M 6-aminocaproic acid, 35% (v/v) glycerol, and 1.0% (v/v) deoxycholic acid. After clarification, the samples were loaded onto 5-13% gradient native gels. Electrophoresis was performed at 100 Volts with 15mA per gel overnight at 4°C. Thyroglobulin (669kDa), Ferritin (440kDa), and catalase (232kDa) (Pharmacia) were used as molecular standards.

GST-PKC ϵ Pull-Down

GST-PKC ϵ affinity pull-down assays were performed as previously reported (142). Briefly, GST-PKC ϵ recombinant protein was generated using the baculovirus system (Pharmingen). Subcellular fractions were incubated with GST-PKC ϵ in binding buffer containing 0.5% (v/v) Triton X-100, 20mM Tris-HCl, pH7.4, 150mM NaCl, 1mM EDTA, 1mM EGTA, and a cocktail of protease inhibitors (Roche) overnight at 4°C. The beads were washed three times and the proteins eluted with glutathione (GSH) elution buffer (Pharmingen).

Sucrose Gradient Purification

Proteins eluted from the GST-PKC ϵ pull-down were overlayed on a 10-30% sucrose gradient (132). Eighteen consecutive fractions, beginning from the top, were collected. Thyroglobulin (669kDa) and catalase (232kDa) were used as molecular standards. Protein fractions were precipitated with cold 10% (w/v) TCA and 90% (v/v) acetone, resolved by SDS-PAGE, and visualized by silver staining. To identify PKC ϵ signaling complexes, the SDS-PAGE was blotted by PKC ϵ antibodies.

Phosphorylation of Akt or eNOS by PKC ϵ

Recombinant Akt or eNOS was incubated with PKC ϵ in PKC reaction buffer containing 0.03mg/ml L α -phosphatidyl-L-serine, 2.5 μ g/ml phorbol 12-myristate 13-acetate, 3.5mM dithiothreitol (DTT), 100 μ M ATP, 6.5mM magnesium chloride, 50mM Tris-HCl, pH7.5 at 30°C for 30min. The reaction was terminated by addition of 5 \times Laemmli sample buffer and boiled 5 min at 90°C. The proteins were then separated by SDS-polyacrylamide gel electrophoresis (PAGE), dried and exposed to X-ray films.

Akt Activity Assay

Akt kinase activity assay was measured using a commercial available kit from Cell Signaling. Briefly, Akt immunoprecipitant were incubated with 1 μ g GSK-3 fusion protein in the presence of

ATP and kinase assay buffer containing 25mM Tris-HCl (pH 7.5), 5mM β -glycerolphosphate, 2mM dithiothreitol, 0.1mM Na_3VO_4 , 10mM MgCl_2 at 30°C for 30 min. Phosphorylation of GSK-3 is measured by Western blotting using a phosphor-GSK-3 (Ser21/9) antibody.

eNOS Activity Assay

eNOS activity was assayed via measuring NO_2/NO_3 production using colorimetric assay kit from Calbiochem. Briefly, incubate recombinant eNOS with/out pretreatment or eNOS immunoprecipitant from mouse heart with reaction buffer containing 1.25mM NADPH, 1.25mM arginine, 0.5mM CaCl_2 , 3 μM tetrahydrobiopterin (BH_4), 1 μM flavin adenine dinucleotide, 1 μM flavin adenine mononucleotide and 25mM Tris-HCl, pH7.4 for 30 min at 30°C. Then nitrate was converted to nitrite with nitrate reductase treatment. Total nitrite was measured by using the Griess reaction as modified by Gilliam et al (68).

Statistical Analysis

Data are reported as mean \pm SEM. Differences among the 4 experimental groups were analyzed using one-way ANOVA. If the ANOVA showed an overall significance, post hoc contrasts were performed with Student t tests for unpaired data using the Bonferroni correction (174).

CHAPTER III

Functional proteomic mapping of the myocardial PKC ϵ signaling subproteome in the mouse heart: a subcellular compartment study

Abstract

Functional proteomic analyses have revealed that protein kinase C ϵ (PKC ϵ), a critical mediator of cardiac protection, forms multiprotein signaling complexes in the myocardium. We previously characterized at least 93 PKC ϵ -associated proteins in the PKC ϵ signaling subproteome including structural proteins, signaling molecules, stress-activated proteins, transcriptional/translational factors, metabolism-related proteins, and PKC-binding domain containing proteins. However, these studies were performed in the context of the whole myocyte, and information regarding the subcellular distribution of these proteins, which is essential for the construction of a cellular map of this subproteome, was lacking. The present studies were designed to investigate the localization and composition of PKC ϵ complexes in each subcellular compartment and the putative mechanisms that govern their assembly. Accordingly, we examined three critical aspects of multiprotein complexes in the myocardium: the molecular size of native protein complexes; the molecular composition of individual protein complexes, and the assembly of each complex in various cellular compartments. Furthermore, the functional significance of these complexes was assessed by delineating their formation in the genesis of a cardioprotective phenotype. Using a comprehensive functional proteomic strategy that combines various protein isolation, separation, and identification techniques, we found that: 1) the cardiac PKC ϵ signaling subproteome is comprised of various multiprotein PKC ϵ complexes; 2) the native size of each PKC ϵ complex differs over a wide range of molecular weight among cytosolic, membrane, nuclear,

mitochondrial, and caveolar compartments; 3) different PKC ϵ complexes appear to have distinct molecular compositions, which vary with subcellular location, suggesting that subcellular location contributes to the identity of PKC ϵ -associating proteins; and 4) the assembly of these complexes is dynamically modulated in the cardioprotected when compared with the normal heart. Taken together, our data represent the first comprehensive analysis of subcellular PKC ϵ protein complexes and demonstrate that formation of these signaling complexes is subcellular compartment-specific.

Introduction

Molecular interactions among proteins serve as essential mechanisms to facilitate the transmission of messages within the cell. Functional proteomic analyses of several signal transduction systems in various cell types have advanced our understanding of how such messages are relayed (91, 97-99, 113, 142, 168, 173). Whereas the majority of previous studies have examined the functional interaction of only limited molecules, proteomic analysis of an entire subproteome enables delineation of complex molecular interactions in both normal and diseased states, providing a more comprehensive picture of how a given signaling network operates. For instance, an investigation by Husi *et al.* identified 77 proteins within the brain NMDA receptor complex, which include cytoskeletal, adaptor, and signaling molecules (98, 99). In a different study, several proteins, ranging from DNA repair to cytoskeletal regulation, were found to undergo expression and/or post-translational alterations following either activation or inhibition of the serine/threonine protein kinase MEK, a key component of the ERK pathway (113).

Recently, our laboratory has used functional proteomics to study myocardial signaling (52, 142). These analyses of the cardiac PKC ϵ signaling subproteome revealed that PKC ϵ forms multimeric signaling complexes with at least 93 proteins, which can be categorized as structural proteins, signaling molecules, stress-responsive proteins, metabolism-related proteins, transcription- and translation-related proteins, and PKC ϵ binding domain-containing proteins (52, 142). Moreover, cardiac protection facilitated by transgenic activation of PKC ϵ was concomitant with altered recruitment and post-translational modification of many protein residents within the PKC ϵ complexes. Together, these studies indicate that a given signaling element, such as PKC ϵ , can interact with and modulate a wide range of proteins, diverse in both structure and biological function.

Recent studies have demonstrated that subcellular compartmentalization is a critical

factor in defining signaling specificity (97, 173). Therefore, the delineation of a detailed cellular map of a signaling system would be of essential importance to our understanding on how this system supports a biological function. Using whole cardiac cell lysates (52, 142), our previous studies identified many molecular participants that comprise the cardiac PKC ϵ signaling subproteome, however, information regarding the subcellular localization of PKC ϵ associated proteins, which is essential to the mapping of this subproteome, is unknown. The formation and activation of specific PKC ϵ signaling complexes at different subcellular compartments has been implicated to have significant functional consequences (12, 172, 173). Characterization of the subcellular distribution of PKC ϵ complexes should provide key insights into the signaling mechanisms underlying the cardiac phenotypes observed in response to PKC ϵ activation.

Accordingly, in the present study, a systematic examination of PKC ϵ complexes was conducted in the cytosolic, nuclear, mitochondrial, membrane, and caveolar subcellular fractions of mouse hearts, with regard to both their native sizes and their protein compositions. Moreover, the dynamic modulation of each subcellular PKC ϵ signaling complex in response to a phenotypic change was also assessed using cardiac protected PKC ϵ transgenic mice that have been previously established in our laboratory (Figure 8). Our data provides critical information to the mapping of the cardioprotective PKC ϵ signaling subproteome in the murine myocardium.

Results

Subcellular fractionation of mouse hearts and isolation of multiprotein PKC ϵ complexes.

Mouse hearts from transgenic PKC ϵ (TG) and nontransgenic (NTG) littermates were homogenized and the lysates were fractionated by differential centrifugation methods into cytosolic, nuclear, mitochondrial, membrane and caveolar fractions. 2D electrophoresis followed by silver staining revealed largely disparate patterns of proteins among various subcellular fractions (Figure 9). To verify the subcellular fractionation processes, immunoblotting for subcellular compartment-specific proteins was carried out for each fraction. The nuclear-specific protein NuMA was only found in the nuclear fraction, whereas the sarcolemmal Na⁺/K⁺ ATPase was primarily localized in the membrane fraction, and the mitochondrial protein prohibitin-1 was mainly concentrated in the mitochondria (Figure 10). Finally, cytosolic separation was assessed using lactate dehydrogenase (LDH) activity, with >98% of the total LDH activity residing in the cytosolic fraction, and only $\leq 0.85\%$ activity present in any of the other cellular fractions (Figure 10). Therefore, our subcellular fractionation procedure was largely efficient with minimum cross-contamination.

We next examined PKC ϵ expression in each subcellular fraction from both NTG and TG hearts by immunoprecipitating PKC ϵ and then immunoblotting for PKC ϵ . Figure 11 shows that PKC ϵ could be efficiently immunoprecipitated from each fraction, which enabled isolation of multiprotein PKC ϵ complexes for further studies in a subcellular-specific manner. As expected the levels of PKC ϵ in every compartment were considerably higher in the TG mouse hearts when compared with the NTG mouse hearts.

Following immunoprecipitation, multiprotein PKC ϵ complexes from different subcellular fractions were eluted and visualized by either 1D SDS-PAGE (Figure 12) or 2D PAGE (Figure 13) followed by silver-staining. Figure 12 illustrates significant differences in PKC ϵ -associated

proteins in the cytosolic, nuclear, mitochondrial, and membrane subcellular compartments. These differences include the number of proteins present within each subcellular PKC ϵ complex as well as the molecular weights of proteins within each subcellular PKC ϵ complex. A similar trend was found following 2D separation, examples of which are shown in Figure 13. Multiprotein PKC ϵ complexes from cytosolic and nuclear fractions displayed distinct 2D electrophoretic patterns of protein spots. Taken together, our results show that each subcellular fraction contains PKC ϵ signaling complexes. Moreover, our data show that the number and size of the proteins within the PKC ϵ complexes differs in each subcellular compartment. The nature of such differences highlights potential subcellular-specific differences in PKC ϵ dependent signal transduction and cellular function.

Delineation of the physical sizes of native multiprotein PKC ϵ complexes.

Having validated the subcellular fractionation and complex isolation procedures, large-scale homogenization and fractionation of NTG mouse hearts were performed. Subcellular protein mixtures from 15 hearts were analyzed by gel filtration chromatography using a Sephacryl S400 column to separate intact protein complexes on the basis of their physical size. The presence of multiprotein PKC ϵ complexes in the different subcellular fractions was then determined by PKC ϵ immunoblotting. Figure 14 shows the chromatographic profiles of PKC ϵ protein complexes in cytosolic, membrane, mitochondrial, and nuclear fractions, respectively. Although a significant amount of PKC ϵ migrated in the form of either a monomer or a dimer (elution volumes 250-280ml), especially in the cytosolic and nuclear fractions, the migration pattern of PKC ϵ at lower elution volumes demonstrated that PKC ϵ associates with many other proteins in different sized complexes. The most striking finding was that the elution profiles of PKC ϵ protein complexes were very different in each subcellular fraction, again emphasizing the heterogeneity of multiprotein PKC ϵ complexes, both in size and composition, among compartments.

Individual fractions separated by gel filtration were further analyzed for PKC ϵ complex formation. After immunoprecipitation with PKC ϵ antibodies, PKC ϵ complex proteins were eluted

and cross-linked with a Sulfo-NHS-LC-Biotin conjugating agent. The biotinylated proteins were then separated by 1D SDS-PAGE and visualized by immunoblotting using a streptavidin-HRP conjugate (Figure 15a). The results confirmed that PKC ϵ associates with a battery of proteins in the mouse heart. Again, it was evident that the collection of proteins within the PKC ϵ complexes differed across individual subcellular compartments. As expected, the larger complexes were found in the lower elution volumes and contained more components in general; whereas the smaller complexes were found in the higher elution volumes and hosted fewer proteins.

To insure that our findings regarding these complexes were not due to protein aggregation artifacts, we performed additional experiments to study these multiprotein complexes using recombinant GST-PKC ϵ as bait proteins. Cardiac subcellular fractions were incubated with GST-PKC ϵ fusion proteins and allowed to form complexes. These complexes were then isolated from the cell lysate by GST pull-down and further fractionated by sucrose gradient centrifugation. Gel analysis and immunoblotting of sucrose gradient fractions showed that in all subcellular compartments (Figure 15b shows cytosolic as an example), GST-PKC ϵ forms specific multiprotein complexes of varying sizes. The peak sedimentation fraction for PKC ϵ appeared to migrate at ~200kDa; however, PKC ϵ containing fractions were also traced at higher weights (>669kDa), showing that large complexes were also formed (Figure 15b). In further complex formation tests, analysis of two selective fractions from gel filtration by deoxycholate-buffered native gel separation definitively show that PKC ϵ is present in several high molecular weight complexes (>880kDa). Combined with the expression profile of each subcellular fractionate generated by gel filtration, these data show that PKC ϵ forms numerous complexes in each compartment that are largely heterogeneous in size and molecular weight (Figure 15c).

Characterization of protein compositions of subcellular PKC ϵ complexes in the mouse heart

In order to assess the identity of the proteins within the subcellular PKC ϵ complexes, PKC ϵ was immunoprecipitated from each fraction and the complexes analyzed by immunoblotting for proteins previously demonstrated to be components of the total cardiac PKC ϵ complex (52, 142).

Protein expression profiles of the PKC ϵ signaling complexes in cytosol

The number of proteins constituting the cytosolic PKC ϵ signaling complex was extensive (Table 1). Many contractile proteins found in the soluble fraction, such as troponin I and actin, resided in the PKC ϵ complex. Interestingly, RACK2 and AKAP79, which are anchoring proteins for several kinases, including PKC, were also associated with cytosolic PKC ϵ complexes. Moreover, many signaling proteins interacted with cytosolic PKC ϵ including lipid kinases (e.g. PI 3-kinase) (Figure 16), tyrosine kinases (e.g. Lck, Pyk2), serine/threonine kinases (e.g. Akt, MAPK), and protein phosphatases (e.g. PP1, PP2A). Stress-activated proteins involved in the synthesis of low molecular weight mediators such as nitric oxide (eNOS) (Figure 16), prostaglandins (COX2), and carbon monoxide (HO-1) were also key components, alongside several members of the heat shock protein family (HSP70, HSP27). The hypoxia sensitive transcription factor Hif-1 α was also found in the cytosolic complex, consistent with the cytoplasmic localization of Hif-1 α in quiescent cells (37), as were multiple metabolic proteins, such as creatine kinase.

Protein expression profiles of the PKC ϵ signaling complexes in nucleus

The nuclear PKC ϵ complex was as extensive in composition as the cytosolic fraction (Table 1). Not surprisingly, the nuclear proteins LAP2, NuMA, and lamin C bound to PKC ϵ in this fraction. A similar profile of signaling kinases and stress-activated proteins that was seen in the cytosol was also observed in the nuclear signaling complex. However, unlike the soluble fraction, we also found the tyrosine kinases Src and BMX in the nuclear PKC ϵ complex. In addition, multiple transcriptional proteins resided in the nuclear PKC ϵ complex, including c-Fos, c-Jun, NF- κ B, Elk-1, CREB, and STATs.

Protein expression profiles of the PKC ϵ signaling complexes in membrane

Consistent with the hypothesis that RACK proteins mediate PKC ϵ translocation to membranous structures, we found both RACK1 and RACK2 in the membrane PKC ϵ complex (Table 1).

AKAP79 also co-localized with PKC ϵ complex in this compartment. Other structural proteins in the membrane complexes included the intermediate filament protein desmin, the tight junction protein villin, the Golgi matrix protein GM130, and β' -adaptin, a constituent of clathrin-coated pits. Along with the many kinases and stress-activated proteins already observed in the cytosolic and nuclear complexes, the cardiac-specific gap junction protein connexin-43 also interacted with PKC ϵ in the membrane.

Protein expression profiles of the PKC ϵ signaling complexes in caveolae

Although the cytosolic, nuclear, and membranous PKC ϵ signaling complexes contained a large number of molecules, we found that the expression profile of proteins in the caveolar PKC ϵ complex was by comparison limited (Table 1). The structural protein caveolin-3 was a major component of the PKC ϵ complex in this specialized micro-domain. RACK2 and β' -adaptin were also found. Again, several kinases were associated with PKC ϵ in caveolae including Ras, and Akt, the tyrosine kinases Src and Lck, and all three MAPK families. In agreement with previous investigations demonstrating strong caveolar localization of NOS (80, 151), both eNOS and iNOS interacted with PKC ϵ in mouse cardiac caveolae (Figure 7).

Protein expression profiles of the PKC ϵ signaling complexes in mitochondria

Our initial analysis revealed number of PKC ϵ associating proteins in the mitochondrial fraction (Table 1). Notably that these proteins include the mitochondrial import protein, prohibitin-1, and the mitochondrial metabolic proteins ANT, VDAC, and isocitrate dehydrogenase. In addition, the signaling kinase profile of the mitochondrial PKC ϵ complex contained multiple tyrosine kinases, as well as members of the MAPK and Akt signaling pathways. There were also a number of stress proteins associating with PKC ϵ in the mitochondria, all pertaining to mediator synthesis and metabolism, such as eNOS (Figure 7) and COX2.

Modulation of subcellular PKC ϵ complexes size in protected myocardium.

It has been previously demonstrated that cardiac-specific transgenic activation of PKC ϵ renders the mouse heart resistant to ischemia/reperfusion injury (136, 142, 145). In addition, analysis of the PKC ϵ complexes from the total cell lysates of protected hearts and wild type mouse hearts showed alterations in the interaction of many proteins with these complexes (52, 142). However, it remained unknown what mechanisms regulated the association of given proteins with PKC ϵ . Thus, the present study investigated the role of subcellular location as a regulatory determinant of the formation of these complexes during phenotypic changes. Gel filtration experiments were performed using PKC ϵ TG mice hearts and then compared to those conducted with NTG mouse hearts to determine the alterations of native subcellular PKC ϵ complexes during cardiac protection. As seen in Figure 6, the native protein profiles of PKC ϵ complexes in TG hearts were also subcellular compartment-specific. In general, the size of native PKC ϵ complexes were increased in the PKC ϵ cardioprotected hearts, with a concomitant decrease in the free forms, that is, the monomeric and dimeric forms, of PKC ϵ , further supporting the concept that protective PKC ϵ signaling involves the recruitment of proteins into these complexes. In particular, in the cytosolic fraction, PKC ϵ formed putatively several very high molecular weight (>MDa) complexes in the TG heart that were not present in the NTG heart. The mitochondrial, membrane, and nuclear PKC ϵ profiles also exhibited dramatic changes in the TG mouse hearts compared with those in NTG hearts. Taken together, these data indicate that PKC ϵ recruits different molecules into the complexes depending on the subcellular location and that the subcellular complexes are individually modified in the protected versus the naïve heart. This finding suggests that this subcellular organization of multiprotein signaling complexes contributes to the simultaneous modulation of multiple functions that lead to a given phenotype, in this case cardiac protection.

In addition to modulation of native PKC ϵ complex size, the composition of each PKC ϵ complex was altered in a subcellular-specific manner in the TG mouse hearts. Indeed, striking variation in the expression profiles of PKC ϵ complexes was observed across different cardiac subcellular locations when normal hearts were compared to those protected against ischemic

injury (Table 2). Enhanced binding of a variety of signaling and stress-activated molecules to PKC ϵ was for the most part subcellular compartment-specific. For example, eNOS was upregulated in all the subcellular PKC ϵ complexes whereas the 170kDa isoform of PI 3-kinase was only increased in the cytosolic and membrane fractions (Figure 16). Regarding structural proteins, the PKC ϵ -selective receptor protein, RACK2 exhibited greater interaction with PKC ϵ in both the cytosol and the membrane fraction of the TG mouse hearts, whereas interaction of PKC ϵ with the LIM/PDZ-containing protein Oracle, a protein that has been characterized in cardiac development (93, 137, 189), was only elevated in the nuclear fraction. The upregulation of these specific proteins in specific subcellular complexes points to the ability of PKC ϵ to modulate multiple, discrete cellular functions.

Discussion

Although several efforts in whole organelle proteomic analyses of mitochondria (12, 77, 118), Golgi (17, 178) and other subcellular fractions (36, 90, 109, 126, 131, 139) have achieved fruitful results, comprehensive characterization of the entire proteome of a particular organelle remains a Herculean task. Major challenges include the dynamic nature of these organelle proteomes with respect to the changes of time and cellular functions. In contrast, studies focusing on the examination of specific subproteomes, such as the PKC ϵ signaling system, will target a sub-group of proteins on the basis of the specific biological functions they perform and will define temporal alterations of these proteins (during the time course of a phenotype change). Consequently, this approach is more focused and will enable us to circumvent the aforementioned challenges. In accord with this concept, simultaneous proteomic analysis of a particular subproteome in all subcellular compartments will illustrate a cellular map of this subproteome, and would offer important novel insights.

Consequently, the primary objectives of the present investigation were to delineate the subcellular distribution of the PKC ϵ signaling subproteome and to understand how assembly of this subproteome is modulated during cardioprotection. We hereby report that the PKC ϵ signaling subproteome is comprised of various multiprotein PKC ϵ signaling complexes; these complexes differ in their native molecular size, and the formation of these complexes is modulated on a subcellular basis within cardiac cells. To our knowledge, this is the first study to demonstrate the formation of native, signaling kinase (PKC ϵ)-based, multiprotein complexes, and to define how modulation of these complexes directly correlates with a specific phenotype (cardioprotection). This study provides three lines of novel information: 1) these data show differential regulation of multiprotein complex assembly across subcellular locations, suggesting that this subcellular organization of distinct multiprotein complexes may serve as a mechanism for signal transduction; 2) these data demonstrate that a signaling kinase, namely PKC ϵ , can form

protein complexes with numerous other functionally diverse proteins, and that the distinct molecular components of these complexes appear to be modulated in a stimulus and subcellular location-specific manner; and most importantly, 3) these findings suggest that analysis of intact, non-denatured multiprotein complexes is an essential step to further our understanding of the manner in which these complexes transduce signals in normal and diseased organs.

We achieved mapping of the PKC ϵ subproteome using two complimentary approaches. The first approach encompasses the biochemical methods of gel filtration, native gels, and 2D electrophoresis, which enabled the determination of the size and composition of the protein complexes. The data presented herein shows clearly that PKC ϵ forms multiprotein complexes that contain many components. Size fractionation of subcellular proteins using gel filtration showed a dynamic size range of PKC ϵ complexes, which differ cross subcellular compartments, and are dynamically modulated in the non-transgenic and cardioprotected PKC ϵ transgenic hearts (Figure 14). Consistent with these findings, GST-PKC ϵ pull-down and sedimentation sucrose gradient analysis illustrate that multiprotein PKC ϵ complexes are heterogeneous and multimeric in nature *in vivo* and *in vitro*. Similarly, the 2D electrophoretic display of PKC ϵ complexes from cytosol and nuclear subcellular fractions yielded different patterns (see Figure 13c) concurrent with the hypothesis that each subcellular fraction contains distinct PKC ϵ complexes.

The second approach taken was the systematic identification of proteins within individual subcellular compartmentalized PKC ϵ complexes, using classical immunoprecipitation and immunoblotting techniques (Table 1 and 2). The majority of previous proteomic studies have relied on mass spectrometry for protein identification. Indeed, using this strategy, our laboratory has previously identified 93 proteins in the PKC ϵ signaling subproteome (52, 142). However, in view of the concerns of the sensitivity of our established mass spectrometry data, we have performed a substantial amount of immunoblotting analyses to verify and substantively confirm these results (52, 142).

The data revealed that the number and nature of specific proteins present within a PKC ϵ complex was dependent upon the subcellular location (see Table 1). For example, the cytosolic

PKC ϵ complexes contained ~50 proteins from all of the functional groups (signaling, structural, stress-activated, transcription, metabolism, and PKC binding domain). In stark contrast, the caveolar PKC ϵ complexes contained only 13 proteins that were primarily signaling and structural molecules; presumably a direct reflection of the specialized nature of this particular subcellular compartment. A somewhat surprising finding was that a large number of PKC ϵ -associated proteins were found outside of their normal subcellular context. The specific PKC ϵ anchor protein RACK2 is thought to localize to cytosolic and membranous compartments such as the sarcolemma and Golgi (41) and indeed co-localized with PKC ϵ in these fractions. However, it was also found to be a key component of the mitochondrial, nuclear, and caveolar complexes. This is an important observation, as it represents a means by which PKC ϵ can localize to these organelles, as has been described during cardiac protection. Similarly, we found that eNOS interacts with PKC ϵ in every compartment (see Figure 16). Many studies have demonstrated the cytosolic and particularly membrane/caveolar localization of eNOS in cardiac cells (24, 180), and thus, the interaction of eNOS with PKC ϵ in these compartments was expected. However, eNOS also co-localized with PKC ϵ in the nuclear and mitochondrial fractions. Nuclear translocation of eNOS in response to VEGF and estradiol (54) has been reported and the formation of nuclear PKC ϵ -eNOS complexes may allow a more direct modulation of transcription by nitric oxide. Nitric oxide is able to permeate the mitochondrial membrane and can influence the function of many components of oxidative phosphorylation (31). Consequently, mitochondrial PKC ϵ -eNOS complexes may enable PKC ϵ to regulate myocardial energy. The functional importance of signaling between PKC ϵ and eNOS has been further elucidated in detail and these findings will be presented in the subsequent Chapter. These examples show that, in addition to further delineating the cardiac PKC ϵ signaling system, the present study provides novel insight into the subcellular expression of a number of molecules and therefore evidence for heretofore unknown functions of PKC ϵ .

The finding that the protein composition of these subcellular PKC ϵ complexes was modulated during cardiac protection serves as important evidence for the role of these complexes

to regulate phenotype. As shown in our previous study (142) many of the PKC ϵ associated proteins have been implicated in protection of the heart against ischemic injury cardioprotection. To name a few, Src (146, 160), Lck (144, 145), Akt (39), MAPKs (12, 141), COX2 (78, 158), NOS (78, 181), connexin-43 (47), NF- κ B (115, 181), HO-1 (184), and HIF-1 (37) have all been shown to play protective roles in the heart. Interestingly, compartment-specific alterations in the association of these known cardiac protective proteins within PKC ϵ complexes were saliently observed in the present study. For example, recruiting NF- κ B to PKC ϵ was only observed in the nuclear fraction whereas increased binding to the caveolin-3 was only found in the caveolar fraction. Again, these data are consistent with the specialized functions of the proteins involved: transcription in the case of NF- κ B and the structure/regulation of caveolae in the case of caveolin-3.

In summary, the present study demonstrates that different subcellular locations within the myocardium contain PKC ϵ complexes with various native sizes and distinct molecular components, suggesting that specific cellular tasks at different subcellular locations are facilitated by regulation of these protein complexes. Furthermore, these subcellular complexes are independently modulated in terms of size and composition when the myocardium is protected against ischemic insult. Taken together, these findings support the notion that subcellular-specific organization of multiprotein complexes may serve as a mechanism by which the cell controls its phenotype.

Figure 8. Flow chart for isolation and visualization of PKC ϵ multiprotein signaling complexes – experimental strategy for Chapter III Mouse hearts were homogenized and separated into different subcellular fractions. Then gel filtration chromatography, PKC ϵ immunoprecipitation or GST-PKC ϵ affinity pull down assay were applied to isolate protein complexes. Different methods were subjected to visualize PKC ϵ multiprotein complexes, leading to obtain various data shown in the following figures.

Tissue Homogenization and
Isolation of Subcellular
Compartments

Isolation / Separation of
Protein Complexes

Visualization of PKC ϵ
Protein Complexes





Figure 9. Two dimensional (2D) electrophoresis separating protein mixtures obtained from mouse heart cytosolic (left) and membrane (right) subcellular fractions. The mouse hearts were homogenized and separated into cytosolic, membrane, mitochondrial and nuclear fractions. Subcellular extracts from cytosolic and membrane fractions were loaded onto immobilized pH gradient (IPG) strips and subjected to isoelectric focusing (pI) in the first dimension. The IPG strips were re-equilibrated and subjected to SDS-PAGE for the second dimension by molecular weight (MW). Proteins were then visualized by silver staining

Cytosolic Fraction



Membrane Fraction




Figure 10. Immunoblotting analysis of known marker proteins in subcellular fractions. Subcellular-specific marker proteins were used to verify the separation procedure, such as nuclear-specific NuMA (upper panel); membrane-specific Na⁺/K⁺ ATPase (middle panel); and mitochondrial-specific Prohibitin-1 (lower panel). Cytosolic-specific lactate dehydrogenase (LDH) activity was also measured in each subcellular fraction (graph).





Figure 11. Characterization of multiprotein PKC ϵ signaling complexes-A. Protein from non-transgenic (NTG) and PKC ϵ transgenic (TG) mouse hearts were immunoprecipitated (IP) with PKC ϵ antibodies, separated by 1D SDS-PAGE, and immunoblotted (IB) with PKC ϵ antibodies. This figure showed that PKC ϵ could be efficiently immunoprecipitated from each subcellular fraction. Moreover, as expected that PKC ϵ protein expression level in every compartment were higher in PKC ϵ transgenic mice when compared with NTG.




Figure 12. Characterization of multiprotein PKC ϵ signaling complexes-B.

Subcellular extracts were subjected to PKC ϵ immunoprecipitation to isolate PKC ϵ protein complexes. Then PKC ϵ associated proteins were separated by 1D SDS-PAGE, and visualized by silver staining. PKC ϵ associated with many proteins, and the patterns of polypeptides were significantly different in each subcellular fraction. The differences were including the number of proteins and protein molecular weight.

PKC ϵ IP : 1D SDS-PAGE : Silver Staining




Figure 13. Characterization of multiprotein PKC ϵ signaling complexes-C.

Subcellular extracts were subjected to PKC ϵ immunoprecipitation to isolate PKC ϵ protein complexes. Then PKC ϵ associated proteins were separated by 2D SDS-PAGE, and visualized by silver staining.

PKC ϵ IP : 2D SDS-PAGE : Silver Staining




Figure 14. Demonstration of PKC ϵ multiprotein signaling complexes by liquid chromatography. Subcellular protein lysates (~10mg) of both non-transgenic (NTG) and PKC ϵ transgenic (TG) hearts were fractionated by gel filtration. Proteins from elution fractions were precipitated and PKC ϵ signaling complexes visualized by immunoblotting. All elution fractions were normalized for PKC ϵ expression using 100 μ g NTG heart lysates as internal control. Bar graph indicates the presence of PKC ϵ multiprotein complexes. The elution profiles of PKC ϵ complexes were different in different subcellular compartments. Compared with non-transgenic animals, the elution profiles of PKC ϵ complexes were shifted toward higher PKC ϵ complexes in PKC ϵ -cardioprotective transgenic mouse hearts.







Figure 15. Determination of the native size of multiprotein PKC ϵ complexes.

(a) The 120th and 180th elution fractions from the gel filtration column were immunoprecipitated with PKC ϵ antibodies. PKC ϵ protein complexes were biotinylated with a trifunctional cross-linking agent (EZ-Link Sulfo-NHS-LC-Biotin [Pierce]). The biotinylated proteins were separated by 1D SDS-PAGE and visualized by immunoblotting using a streptavidin-HRP conjugate. (b) Native PAGE analysis of 180th elution fraction. Left column: various cellular multiprotein complexes visualized by native PAGE followed by silver staining. Right column: the PKC ϵ protein complexes visualized by native PAGE followed by immunoblotting (IB) for PKC ϵ (indicated by arrows). (c) Sedimentation sucrose gradient analysis of GST-PKC ϵ -associated proteins. GST-PKC ϵ fusion proteins were used for the affinity pull down PKC ϵ complexes from cytosol. Subsequently, various sizes of GST-isolated protein mixtures were separated by sucrose gradient (10-30%). Upper panel: PKC ϵ immunoblotting of the sucrose gradient fractions collected (1-to-18), arrows indicating the molecular weight range of PKC ϵ multiprotein complexes (upper arrows at 232kDa and 669kDa, respectively). Lower panel: sucrose gradient fractions (1-to-18) were further separated by SDS PAGE and proteins visualized by silver staining.






Figure 16. Identification of the molecular components of the multiprotein PKC ϵ complexes: examples of eNOS and p170 PI 3-Kinase.

Proteins from non-transgenic (NTG) and PKC ϵ transgenic (TG) mouse hearts were separated into cytosolic, membrane, mitochondrial, nuclear and caveolae fractions. Then subcellular extracts were immunoprecipitated (IP) using PKC ϵ antibodies, separated by 1D SDS-PAGE and immunoblotted (IB) with eNOS antibodies (left panels) and p170 PI 3-kinase antibodies (right panels).





Table 1 : PKCε signaling complex proteins in subcellular fractions

	M _r (kDa)	pI	Known Function	Subcellular Fractionations				
				C	N	M	Mit	Cav
Structural proteins								
Troponin I	24	9.6	Contractile protein	+				
Troponin T	35	5.2	Contractile protein	+				
Desmin	53	5.2	Cytoskeletal protein	+		+		
RACK1	36	7.6	PKC binding protein		+	+		
RACK2	102	5.2	PKC binding protein	+	+	+	+	+
Villin	95	5.7	Cytoskeletal protein	+		+		
β'-Adaptin	106	5.0	Intracellular transport adaptor protein	+		+		+
AKAP79	79	5.1	Anchoring protein	+	+	+	+	
Vimentin	54	5.1	Intermediate filament	+	+	+		
Cadherin-2	129	4.9	Cell junction adhesions, Ca ⁺⁺ regulated	+	+	+	+	
Lamin C	65	6.4	Nuclear intermediate filament	+	+	+		
Oracle1/2	70/80	8.0	Cardiac development. LIM/PDZ domains	+	+	+	+	
Vinculin	117	5.7	Focal adhesion protein	+		+		
Glucose-regulated protein	75	5.1	Molecular chaperone	+		+		
Alpha-Actinin-2	104	5.4	Intermediate filament, Z band	+		+	+	
Lap2	75	8.6	Nuclear structural protein		+			
NMA	240	5.	Nuclear protein involved in mitosis	+	+			
GM130	130	5.0	Golgi matrix protein			+		
Prohibitin-1	30	5.6	Mitochondrial chaperone protein				+	
Caveolin-3	18	5.5	Cardiac caveolae structural protein					+
Signaling proteins								
PI 3-kinase	170	8.3	Phosphoinositide lipid kinase	+	+	+		
Pyk2	116	5.9	FAK tyrosine kinase family	+	+	+		
Lck	56	5.2	Src family tyrosine kinase	+	+	+	+	
Src	60	7.9	Src family tyrosine kinase	+	+	+	+	+
PKB/Akt	59	5.6	Stress-activated kinase	+	+	+	+	+
p38 MAPK	38	5.5	Stress-activated kinase	+	+	+	+	+
JNK1/2	46/54	7.6/6.2	Stress-activated kinase	+	+	+	+	+
ERK1/2	44/42	6.0/8.9	Receptor and stress-activated kinase	+	+	+	+	+
MEK1	45	6.2	Upstream activator of ERK	+			+	
Ras	21	6.2	GTPase that activates MAPK pathways	+	+	+	+	+
Connexin-43	43	9.0	Cardiac gap junction protein	+		+		
PDK1	63	6.7	Ser/Thr kinase	+	+	+	+	
PP2A	36	5.3	Ser/Thr kinase phosphatase	+		+	+	
PP1	38	5.9	Ser/Thr kinase phosphatase	+	+	+		

PI-3-kinase	85	6.0	Phosphoinositide lipid kinase	+	+	+	+
BMX	80	8.6	Tec family tyrosine kinase	+	+	+	
Stress-activated proteins							
iNOS	130	7.8	Inducible nitric oxide synthase	+	+	+	+
eNOS	140	6.6	Constitutive nitric oxide synthase	+	+	+	+
COX-2	70	7.0	Prostaglandin H synthase	+	+	+	
Heme-oxygenase 1	32	6.1	Stress-activated protein	+	+	+	
Aldose reductase	36	6.7	Osmotic stress-activated protein	+	+		+
Heat shock protein 70	70	5.5	Molecular chaperone	+	+	+	+
Heat shock protein 27	25	6.4	Molecular chaperone	+	+	+	
α B-cystallin	20	7.3	Molecular chaperone	+	+	+	
Heat shock protein 60	61	6.3	Molecular chaperone	+			+
Metabolism-related proteins							
ANT1/2	33	9.7	ATP/ADP exchanger	+			+
Creatine kinase	43	6.6	Reversibly phosphorylates creatine, energy storage	+			
GAPDH	36	8.4	Glycolysis	+			
Isocitrate DH	38	6.0	Citric acid cycle	+	+		+
VDAC	31	8.6	Mitochondrial pore component (ion transport)	+			+
Transcription- & Translation-Related Proteins							
Histone H4	14	11.3	DNA binding to facilitate chromatin structure	+	+		
hnRNP H1	49	5.9	mRNA processing and regulation				+
hnRNP K	51	5.3	mRNA processing and regulation	+	+		
Ribonucleoprotein	37	6.4	mRNA processing and regulation, heat shock, splicing	+	+	+	
NF κ B p65	65	5.5	Activated by stress and inflammation	+	+		
HIF-1 α	100	5.2	Hypoxia-inducible transcription factor	+	+		
c-Fos	55	4.8	Proto-oncogene, part of AP1 complex			+	
c-Jun	36	8.9	Proto-oncogene, part of AP1 complex			+	
CREB	43	5.4	Binds cAMP response element			+	
Elk	45	6.3	Activated by MAPK family kinases			+	
STAT1	87	5.4	Activated by JAK family kinases			+	
STAT3	88	5.9	Activated by JAK family kinases	+	+		
c-Myc	50	5.5	Proto-oncogene	+	+		
hnRNP a2/b1	37	9.0	mRNA processing and regulation			+	

All pI values were estimated using the ExPASy compute pI/MW tool available on the web.

C=Cytosol; N=Nuclei.; M=Membrane; Mit=Mitochondria; Cav=Caveolae.

Table 2 : Altered protein expression in PKC ϵ signaling complex proteins in subcellular fractions

	Subcellular Fractionations				
	Cytosol	Nuclear	Membrane	Mitochondria	Caveolae
Structural proteins					
RACK2	++		++		
Villin			++		
AKAP79		++			
Oracle1/2		++			
GM130			++		
Caveolin-3					++
Signaling proteins					
PI 3-kinase (85kDa)					++
Pyk2 kinase	++				
Lck	++		++	++	
Src			++		
PKB/Akt		++	++		
p38 MAPK				++	
JNK1/2		++		++	
ERK1/2			++	++	
Ras	++	+++	+++		
PDK1			++	++	
BMX		++	++	++	
PI 3-kinase (170kDa)	++		++		
Stress-activated proteins					
eNOS	++	++	++	++	++
Aldose reductase	++			++	
Heat shock protein 27	++	++			
α B-cystallin			++		
Transcription- & Translation-Related Proteins					
NF κ B p65		++			
HIF-1 α		++			
c-Jun		++			
STAT3		++			

The “++” signs indicate increased expression of proteins in various subcellular compartments in cardiac protected PKC ϵ transgenic mice. “++” reflects a change of 1-2 fold, “+++” a change of 2-3 fold.

CHAPTER VI

Functional proteomic analysis of myocardial PKC ϵ -Akt-eNOS signaling modules during cardioprotection

Abstract

Having established in the studies described in the foregoing Chapter that PKC ϵ complexes are differentially regulated in a subcellular compartment specific fashion, we next wanted to determine the manner in which individual signal transduction modules are formed to transducer signaling within these complexes.

Among the 93 different PKC ϵ -associated proteins that we identified to date, the protective kinase Akt and stress-activated protein endothelial nitric oxide synthase (eNOS) are of importance due to their independent abilities to promote cell survival and prevent cell death. Their co-localization within the PKC ϵ protein complex suggests that PKC ϵ , Akt and eNOS may form signaling modules for the regulation of nitric oxide (NO) production and hence the manifestation of the cardioprotective phenotype. Accordingly, using functional proteomic tools, we found that: (1) PKC ϵ , Akt and eNOS interact and form signaling modules both *in vitro* and *in vivo*; activation of either PKC ϵ or Akt enhances the formation of PKC ϵ -Akt-eNOS signaling modules; moreover, the formation of these signaling modules is dynamically regulated during cardioprotection; (2) PKC ϵ directly phosphorylates and enhances activation of Akt *in vitro*, and PKC ϵ transgenesis increases both phosphorylation and activation of Akt *in vivo*; (3) PKC ϵ can also directly phosphorylate eNOS *in vitro*, and this phosphorylation enhances eNOS activity. Furthermore, overexpression of PKC ϵ *in vivo* increased phosphorylation of eNOS at Ser1177, representing increased eNOS activity. This study characterizes, for the first time, functional as well as physical

coupling of PKC ϵ , Akt, and eNOS in the heart, and provides novel information regarding the formation and modulation of PKC ϵ -Akt-eNOS signaling modules during cardioprotection.

Introduction

The proteome is proposed to be composed of multiple integrated functional subproteomes, each of which is organized to facilitate a unique function. Moreover, specific signaling modules are formed within these subproteomes in response to different stimuli (82, 99, 170). In particular, our laboratory has focused on the PKC ϵ subproteome, and the modules present within. PKC ϵ plays a critical role in the genesis of cardioprotection against myocardial ischemic injury (49, 75, 140, 147), and cardiac-specific expression of PKC ϵ is cardioprotective (142). Recent studies in our laboratory have found that PKC ϵ forms multimeric signaling complexes with at least 93 proteins, including structural proteins, stress-activated proteins, proteins associated with transcription and translation, proteins involved in cellular metabolism, and various kinases and phosphatases (52, 142, 160, 172). Furthermore, the assembly of these multiprotein PKC ϵ complexes is dynamically modulated in PKC ϵ -transgenic mouse hearts, which serves as important evidence for the role of these complexes in regulating the cardioprotective phenotype. Among the 93 known protein partners, Akt and eNOS are of great interest as they are well established to independently promote cell survival and prevent cell death in response to various forms of stress (16, 42, 63, 182).

Previous studies in non-cardiac cells demonstrate physical and functional interactions among PKC, Akt and eNOS. Akt can directly interact and phosphorylate eNOS on Ser1177 (63). Akt co-immunoprecipitates with PKC ζ or PKC ι (48, 119), and the PH domain of Akt has been implicated as the binding site for PKC ζ to Akt1 and Akt3 (110). Li et al show that overexpression of PKC α stimulates Akt activity and suppresses apoptosis induced by interleukin 3 withdrawals in myeloid progenitor cells (114). In contrast, Wen and colleagues (176) report that in A549 and HEK293 cells, the phosphatidylinositol 3-kinase/Akt signaling pathway is negatively regulated by PKC. Moreover, it is reported that activation of PKC inhibited eNOS activity by attenuating eNOS

phosphorylation on Ser1177 and increasing phosphorylation of Thr495 in endothelial cells (123, 124). However, despite this information nothing is known regarding the formation and regulation of PKC ϵ -Akt-eNOS signaling modules in myocytes, especially during cardioprotection. Therefore, a comprehensive investigation needs to be implemented to draw a clearer picture of the temporal and spatial relationship among PKC ϵ , Akt and eNOS in the regulation of NO production and the genesis of cardioprotection.

Accordingly, we hypothesized that PKC ϵ , Akt and eNOS form a signaling module within the mouse heart, the assembly and activation of which contributes to the genesis of cardioprotection. By applying functional proteomic tools, we examined the PKC ϵ -Akt-eNOS signaling modules *in vitro* and *in vivo* in two major aspects: molecular architecture, as defined by protein-protein interactions; and module regulation, as defined by posttranslational modification and alteration of activity of each component of PKC ϵ -Akt-eNOS signaling. Moreover, using mice with cardiac-specific expression of active PKC ϵ , we also assessed how these signaling modules are regulated during cardioprotection.

Results

PKC ϵ can directly interact with Akt and eNOS *in vitro*

To examine the ability of PKC ϵ to directly interact with Akt or eNOS, GST-PKC ϵ fusion proteins were incubated with recombinant Akt or eNOS, followed by SDS-PAGE and Western blotting for Akt or eNOS. Figure 17 demonstrates that PKC ϵ directly interacts with Akt *in vitro*. Moreover, we also found that GST-PKC ϵ preferentially interacts with recombinant active Akt that is expressed by baculovirus in Sf21 insect cells and activated with GST-MAPKAPK2 and PDK1, as compared to a much weaker interaction with inactive Akt from insect cells without treatment. Furthermore, we assessed the domain-specificity of the interaction between Akt and PKC ϵ by incubating GST-PKC ϵ with recombinant Akt protein with or without PH domain. We found that the interaction between GST-PKC ϵ and mutant, PH domain deficient, Akt was significantly attenuated, suggesting that this region of Akt is responsible for its interaction with PKC ϵ (Figure 17a and b). Taken together, our data indicated that PKC ϵ directly interacts with Akt primarily through binding Akt's PH domain. Using a similar approach, we also found that GST-PKC ϵ could also physically directly interact with eNOS, as shown in Figure 17c.

Since PKC ϵ can directly interact with Akt and eNOS *in vitro* in a binary sense, we then assessed the ability of these molecules to enhance their association with each other beyond that seen in the foregoing pair-wise experiments. Recombinant Akt and eNOS were incubated with or without PKC ϵ followed by eNOS immunoprecipitation and Akt immunoblotting. We found that recruitment of PKC ϵ increased the binding affinity of eNOS with Akt (Figure 18a left). To examine the effect of PKC ϵ activity on this interaction, recombinant PKC ϵ was next incubated with Akt and eNOS in the presence or absence of potent PKC activators phorbol ester (PMA) and phosphatidylserine (PS) followed by immunoprecipitation of PKC ϵ and blotting for Akt or eNOS.

Figure 18a (right) showed that although PKC ϵ formed protein complexes with Akt and eNOS in the absence of PKC activators, PMA/PS-activated PKC ϵ exhibited a much high affinity for Akt and eNOS, indicating that activation of PKC ϵ enhanced its interaction with both Akt and eNOS. In addition, we tested the effect of Akt on the interaction of PKC ϵ and eNOS by incubating PKC ϵ and eNOS with active or inactive Akt recombinant proteins. Similar to the results obtained for PKC ϵ , activation of Akt was found to enhance its interaction with PKC ϵ and eNOS, and its ability to facilitate interactions between PKC ϵ and eNOS (Figure 18b).

PKC ϵ interacts with Akt and eNOS *in vivo*

Previous studies have shown that Akt and eNOS are present in cardiac PKC ϵ -signaling complexes, as assessed by both GST-PKC ϵ affinity pull down assay and PKC ϵ immunoprecipitation (52, 142), indicating that PKC ϵ can interact with Akt and eNOS *in vivo*. In the present study, using PKC ϵ immunoprecipitation, we found that although all three isoforms of Akt are present in the mouse heart, only Akt1 and Akt2 interact with PKC ϵ (Data not shown).

To assess the formation of PKC ϵ -Akt-eNOS signaling modules *in vivo*, mouse hearts were homogenized and subjected to gel filtration chromatography using a Sephacryl S400 column. This process separates intact myocardial protein complexes on the basis of their physical size. Individual fractions from liquid chromatography were next immunoprecipitated with PKC ϵ antibodies. The PKC ϵ immuno-complexes were then separated by SDS-PAGE and blotted for PKC ϵ , Akt and eNOS. Figure 19a shows the chromatographic profiles of PKC ϵ -associated Akt and eNOS protein complexes in normal mouse hearts. We found that Akt and eNOS co-precipitated with PKC ϵ in multiple fractions, indicating that PKC ϵ , Akt and eNOS interacted with each other in a variety of different sized PKC ϵ -containing complexes. Furthermore, compared with the profiles of non-transgenic mouse hearts, we found that the expression profiles of PKC ϵ -Akt-eNOS signaling modules in PKC ϵ -cardioprotective transgenic mice were shifted towards higher molecular weight complexes (Figure 19b), indicating the dynamic modulation of PKC ϵ -Akt-eNOS signaling modules *in vivo* during cardioprotection.

PKC ϵ can directly phosphorylate Akt and eNOS and enhance their activities *in vitro*.

Based on the above findings, we further investigated the effect of PKC ϵ activation on the formation of PKC ϵ -Akt-eNOS modules *in vitro*. First, we assessed whether PKC ϵ could directly phosphorylate Akt and/or eNOS *in vitro*. Recombinant Akt or eNOS were incubated with recombinant PKC ϵ in the presence of the PKC activators, phorbol ester (PMA) and phosphatidylserine (PS), and radiolabeled P³²- γ -ATP. After SDS-PAGE separation, phosphorylation was visualized by autoradiography. We found that PKC ϵ could directly phosphorylate both Akt and eNOS (Figure 20a).

Next, we further tested the effects of these PKC ϵ -induced phosphorylation modifications on the activation status of Akt and eNOS. After incubating recombinant inactive Akt with PKC ϵ , Akt activity was assessed by its ability to phosphorylate GSK-3 β . We found that Akt activation was dramatically increased with the PKC ϵ treatment (Figure 20b) as compared with inactive Akt alone, indicating that activated PKC ϵ dramatically enhances Akt activity *in vitro*. Similarly, we found that PKC ϵ treatment significantly enhanced eNOS activity, as measured by nitrate/nitrite production, (178.8 \pm 11%, P<0.05 vs eNOS only) when compared with non-treatment (Figure 20b).

Studies by other investigators indicate that Akt is activated by phosphorylation of Thr308 within its activation loop which serves to stabilize the molecule in its active confirmation. Subsequently, phosphorylation of serine 473 (Ser473) at the C-terminus is essential for full activation of Akt. While it is well known that PDK1 is responsible for Thr308 phosphorylation, the upstream kinases that catalyze the phosphorylation of Ser473 is unclear. Because our initial studies demonstrated signal transduction between PKC ϵ and Akt, we next tested whether PKC ϵ could be a putative PDK2 that targets Ser473 by using the site-specific phospho-antibody of Akt after the PKC ϵ treatment. Figure 21 shows that addition of recombinant PKC ϵ significantly increased Ser473 phosphorylation on Akt as compared with that of inactive Akt alone, indicating that *in vitro*, PKC ϵ directly phosphorylates Akt on Ser473.

PKC ϵ transgenesis enhances phosphorylation and activation of Akt and eNOS *in vivo*

In addition to the *in vitro* phosphorylation experiments, we also examined the phosphorylation of Akt and eNOS *in vivo*. PKC ϵ -cardioprotected transgenic mice hearts were used to test the effect of activation of PKC ϵ on Akt by examining the two conserved phosphorylation sites of Akt. Using the site-specific phospho-antibodies of Akt, we found that phosphorylation of Akt on both activation sites Thr308 ($336.1 \pm 15.1\%$ vs NTG) and Ser473 ($181.3 \pm 18.2\%$ vs NTG), was significantly increased in PKC ϵ transgenic mice when compared with non-transgenic animals. These post-translational modification changes occurred in the absence of a total change in Akt protein expression (Figure 22a). Interestingly, total cardiac cellular Akt activity was increased in PKC ϵ cardioprotected mice (Figure 22b top panel). Moreover, when we examined PKC ϵ -associated Akt activity (PKC ϵ complexes were isolated by immunoprecipitation and Akt kinase activity assays performed), we found a surprising augmentation of Akt activity, considerably greater than that seen in the total lysate pool, when Akt was associated with PKC ϵ (Figure 22b lower panel). These data support the concept that module formation between PKC ϵ and Akt accomplishes signal transduction by post-translationally modifying Akt and increasing its activity on downstream substrates.

It is well documented that once Akt is activated, it can phosphorylate eNOS at serine 1177(Ser1177) resulting in activation of eNOS and the production of nitric oxide. Therefore, we examined the phosphorylation of eNOS *in vivo* in the PKC ϵ transgenic mice hearts. Phosphorylation of eNOS on Ser1177 ($407.8 \pm 43.7\%$ vs NTG) was enhanced in the PKC ϵ transgenic mice (Figure 23), indicating that overexpression of PKC ϵ increased eNOS activity *in vivo*. Meanwhile, total eNOS protein expression was also slightly increased ($140.9 \pm 7.2\%$ vs NTG) in PKC ϵ transgenic mice.

Discussion

Recent studies suggest that cellular biological functions, such as signal transduction, are accomplished by multi-protein complexes, which contain within them modules that are composed of several species of interacting molecules. Modules may have discrete functions engendered from the interactions among their individual components. Understanding how these molecules interact and regulate each other is critical in order to comprehend the specific biological function of such modules. Accordingly, in this study, we hypothesized that PKC ϵ forms functional signaling modules with Akt and eNOS, and that the assembly and activation of these modules contributes to the genesis of cardioprotection. Using functional proteomic approaches, we tested the hypothesis in two different aspects: molecular architecture and module regulation. There are several important findings in this investigation. First, PKC ϵ interacts with Akt and eNOS both *in vitro* and *in vivo*, and these interactions are promoted by enhanced activation of PKC ϵ . Second, PKC ϵ forms various sizes of signaling modules with Akt and eNOS, which are dynamically regulated in PKC ϵ transgenic mice during cardioprotection. Third, PKC ϵ directly phosphorylates Akt and eNOS, resulting in increased Akt and eNOS activation *in vitro*, whereas *in vivo*, overexpression of PKC ϵ increases both phosphorylation and activation of Akt and eNOS, especially PKC ϵ -associated Akt activation. This study characterizes, for the first time, physical and functional coupling of PKC ϵ , Akt, and eNOS in the mouse heart, and provides novel information regarding the formation and modulation of PKC ϵ -Akt-eNOS signaling modules during cardioprotection.

Spatial distribution of PKC ϵ -Akt-eNOS signaling modules

In addition to the molecular architecture of signaling modules and their regulation, the spatial distribution of a module within the cell is another important aspect to understand the

function of these signaling units. Using classical immunoprecipitation and immunoblotting techniques, our previous studies already determined the subcellular distributions of PKC ϵ -Akt-eNOS signaling modules both in normal and cardioprotective hearts. We found that, accompanying the increased protein expression of PKC ϵ in multiple fractions (cytosol, membrane, mitochondria, nuclear, and caveolae), protein expression of eNOS in PKC ϵ complexes was also increased in all five subcellular fractions. In contrast, Akt expression was only found to be increased in PKC ϵ -associated *membrane* and *nuclear* fractions in cardioprotected mouse hearts. Therefore, although PKC ϵ -Akt-eNOS modules were found to be existent in multiple fractions, increased formation of this module occurred only in membrane and nuclear fractions during cardioprotection, implicating subcellular compartment-specific functions of PKC ϵ -Akt-eNOS modules in normal and protected hearts. For example, many studies have demonstrated that translocation to the membrane is an important regulatory action for activated PKC ϵ and Akt. Moreover, eNOS is largely membrane-associated as a result of the N-terminal myristoylation and palmitoylation of this molecule (30). Thus, we expected to see the increased formation of PKC ϵ -Akt-eNOS signaling modules in membrane fractions during cardioprotection. Increased formation of PKC ϵ -Akt-eNOS signaling modules in nuclear fractions in PKC ϵ transgenic mice may modify activation of transcription factors or other targets within nucleus fraction and thereby regulate cardioprotective gene/protein synthesis.

PKC ϵ forms different sizes of signaling modules with Akt and eNOS

Revealing the existence of intact, non-denatured, multiprotein PKC ϵ -Akt-eNOS modules is an important step to confirm our hypothesis regarding the involvement of these modules in signal transduction. Using biochemical methods of gel filtration, PKC ϵ immunoprecipitation, we have delineated the physical characteristics of PKC ϵ -Akt-eNOS signaling modules. Our data clearly show that PKC ϵ co-exists with Akt and eNOS in protein complexes of varying sizes in the mouse heart. Moreover, during PKC ϵ -induced cardioprotection, the profile of PKC ϵ -Akt-eNOS modules is dynamically regulated. These data provide PKC ϵ -Akt-eNOS modules shifting to larger

molecular weight complexes. These data provide a solid base for our future study to explore the composition and interaction of each PKC ϵ -Akt-eNOS signaling module.

To confirm the presence of PKC ϵ -Akt-eNOS signaling modules *in vivo*, in addition to separation protein complexes following PKC ϵ immunoprecipitation, we used GST-PKC ϵ pull down assays to form new PKC ϵ complexes, and then separated the complexes by deoxycholate-buffered native gel electrophoresis followed by Akt and eNOS immunoblotting. It is found that only a few of PKC ϵ -Akt-eNOS modules had been formed in non-transgenic mouse hearts, whereas, a high molecular weight PKC ϵ -Akt-eNOS signaling module was found in the PKC ϵ -cardioprotected transgenic mice. The size of this macromolecular complex was estimated to be greater than 880kDa, as determined by comparison to protein markers (data not show). Interestingly, we found significantly fewer PKC ϵ -Akt-eNOS modules via the native PAGE approach as compared with the results from the gel filtration technique. The difference between these results is most likely due to the fact that addition of exogenous PKC ϵ will likely disrupt endogenous preexisting complexes while also inducing formation of other complexes. Thus, these techniques (immunoprecipitation and GST-based affinity pull-down) appear to be complementary, in that they identify non-overlapping pools of protein interactions.

Since PKC ϵ interacts with Akt and eNOS to form varying sizes of complexes, how do these three molecules interact? Several investigators have shown that PH domain is the preferred interaction domain for PKC binding to Akt. In our study, we found that PKC ϵ preferred to interact with Akt via its PH domain which was congruous with previous studies in non-cardiac cells (18). Moreover, we found that PH domain of Akt may not necessary for eNOS binding to Akt *in vitro* (data not shown). However, the detailed interactions among PKC ϵ , Akt and eNOS are still under investigation.

PKC ϵ is a positive regulator for the formation of PKC ϵ -Akt-eNOS signaling modules.

Previous studies have focused on the *individual roles* of PKC ϵ , Akt and eNOS as cardioprotective proteins. These studies include one in which cardiac-specific expression of constitutively active Akt was found to reduce infarct size and apoptosis after ischemia-reperfusion

injury (62). In another investigation, myocardial overexpression of eNOS was found to greatly stimulate cardiac nitric oxide production and to protect the heart against ischemia/reperfusion injury. Despite this, the effect of PKC ϵ on the regulation of Akt and eNOS is controversial, and data regarding the association of these molecules is particularly lacking in cardiac cells. Amino acid sequence analyses reveal that both Akt and eNOS contain multiple potential PKC phosphorylation sites. In our study, we found that PKC ϵ could directly phosphorylate Akt and eNOS *in vitro*, whereas, cardiac-specific overexpression of PKC ϵ induces increased phosphorylation of both Akt and eNOS.

Previous studies have shown that overexpression of PKC α stimulated the activation of Akt. In contrast, several other investigators reported that in A549 and HEK293 cells, the phosphatidylinositol 3-kinase/Akt signaling pathway was negatively regulated by PKC (176). PMA-induced apoptosis was accompanied by inhibition of Akt activity (149). Similarly, PKC activation inhibited eNOS activity by attenuating eNOS phosphorylation on Ser1177 and increasing phosphorylation of Thr495 in cardiovascular endothelial cells (123, 124). Thus, it was unclear in what manner PKC ϵ , Akt and eNOS might interact to regulate each other in the setting of cardioprotection. In the present study, cardiac PKC ϵ was found to serve as a positive regulator of the formation and regulation of PKC ϵ -Akt-eNOS signaling modules in mouse hearts. We found that activation of PKC ϵ enhances the module formation, and the phosphorylation modification induced by PKC ϵ may alter the conformation of both Akt and eNOS and thereby further enhance their activity both *in vitro* and *in vivo*. These data indicate that through the increased formation of PKC ϵ -Akt-eNOS signaling modules, activation of PKC ϵ may promote cell survival and stimulate nitric oxide synthesis, and therefore contribute to the genesis of cardioprotection against ischemic injury.

Regarding the finding that PKC ϵ can directly phosphorylate eNOS and increase eNOS activity *in vitro*, the possibility exists that the purity of eNOS recombinant protein may influence the *in vitro* results. Nonetheless, the increased eNOS activity appear to be PKC ϵ -dependent.



PKC ϵ is a candidate for PDK2

Akt has two conserved phosphorylation sites, Thr308, and Ser473, which are necessary for its maximum activation. Generally, the upstream kinase that fulfils the phosphorylation of Thr308 is PDK-1. But the upstream kinases that are responsible for the Ser473 phosphorylation is unknown, and has been tentatively named "PDK-2". Several candidates have been proposed, including PKC ζ (34, 190), a lipid-raft-associated kinase (84), or simply the autophosphorylation processes followed by the PDK-1 activity (1). In the present study, we tested the hypothesis that PKC ϵ may exhibit PDK-2 activity by phosphorylating Akt Ser473. Interestingly, we found that PKC ϵ did indeed phosphorylate Akt Ser473 *in vitro*, indicating PKC ϵ maybe one of the candidates *in vivo* to target Akt and increase Akt activation by modifying Ser473 phosphorylation. Determination of the ability of PKC ϵ to phosphorylate Akt on Ser473 *in vivo*, and the physiological importance of this post-translational modification is an extensive process that is beyond the scope of the current investigation.

In summary, this study successfully characterized myocardial PKC ϵ -Akt-eNOS signaling modules, which can serve as a good model for future research addressing the signaling mechanisms underlying PKC ϵ -mediated cardioprotection and other phenotypes. The present study provided new information about the molecular architecture of PKC ϵ -Akt-eNOS signaling modules, the subcellular localization of these modules, and the dynamic modulation of these modules during the development of the cardioprotective phenotype. These studies are critical steps towards our goal of defining the cardiac proteome, the multiple integrated functional subproteomes that compose this proteome, and the specific signaling modules required for unique cellular functions.

Figure 17. PKC ϵ directly interacts with Akt and eNOS *in vitro* GST-PKC ϵ fusion proteins were incubated with recombinant Akt or eNOS, followed by SDS-PAGE and Western blotting for Akt or eNOS. a. GST-PKC ϵ interacts with Akt *in vitro*, and preferentially interacts with recombinant active Akt when compared with recombinant inactive Akt; b. GST-PKC ϵ had higher binding affinity with the Akt PH domain when compared with PH domain deleted Akt; c. GST-PKC ϵ also physically interacts with eNOS *in vitro*.

a.



b.





Figure 18. Activation of both PKC ϵ and Akt enhanced PKC ϵ -Akt-eNOS module formation *in vitro*. a. Left, recombinant Akt and eNOS were incubated with or without PKC ϵ followed by eNOS immunoprecipitation and Akt immunoblotting. Recruitment of PKC ϵ increased the binding affinity of eNOS with Akt; right, recombinant PKC ϵ was further incubated with Akt and eNOS plus or minus potent PKC activators phorbol ester (PMA) and phosphatidylserine (PS) followed by immunoprecipitation of PKC ϵ and blotting for Akt or eNOS. PMA/PS-activated PKC ϵ exhibited higher affinity for Akt and eNOS compared with that of PKC ϵ without stimulation. b PKC ϵ and eNOS with active or inactive Akt recombinant proteins. Activation of Akt also enhanced the interaction of PKC ϵ with Akt and eNOS.

a.



b.





Figure 19. Formation of PKC ϵ -Akt-eNOS signaling modules *in vivo*. Mouse hearts were homogenized and subjected to gel filtration chromatography using a Sephacryl S400 column followed by immunoprecipitation with PKC ϵ antibodies and immunoblotting for PKC ϵ , Akt and eNOS. a. The chromatographic profiles of PKC ϵ -associated Akt and eNOS protein complexes in the normal mouse hearts; b the chromatographic profiles of PKC ϵ -associated Akt and eNOS protein complexes in the PKC ϵ -transgenic mouse hearts. Akt and eNOS co-precipitated with PKC ϵ in multiple fractions to form a variety of different sized PKC ϵ complexes. Compared with the profiles of non-transgenic mouse hearts, the expression profiles of PKC ϵ -Akt-eNOS signaling modules in PKC ϵ -cardioprotective transgenic mice were shifted towards higher molecular weight complexes.

a. NTG mouse hearts



b. PKC ϵ TG mouse hearts






Figure 20. PKC ϵ phosphorylates and enhances activation of Akt and eNOS *in vitro*.

a. Recombinant Akt or eNOS incubated with recombinant PKC ϵ in the presence of the PKC activators phorbol ester (PMA) and phosphatidylserine (PS) and P³²- γ -ATP. After SDS-PAGE separation, phosphorylation was visualized by autoradiography; b. After incubating recombinant inactive Akt or eNOS with PKC ϵ , Akt and eNOS activity were assessed by its ability to phosphorylate GSK-3 β or produce nitrate/nitrite, respectively.

a. Akt and eNOS phosphorylation:



b. Akt *In vitro* activity:



eNOS *In vitro* activity:




Figure 21. PKC ϵ directly phosphorylated Akt at its Ser473 site *in vitro*. After incubating PKC ϵ with recombinant Akt, Ser473 phosphorylation was detected by site-specific antibody phospho-Akt-Ser⁴⁷³. Addition of recombinant PKC ϵ significantly increased Ser473 phosphorylation on Akt as compared with that of inactive Akt alone.






Figure 22. Effect of PKC ϵ on the phosphorylation and activation of Akt *in vivo*

a. Non-transgenic and PKC ϵ transgenic mouse hearts were homogenized, and separated with SDS-PAGE followed by immunoblotting with Akt, PKC ϵ and site-specific phospho-Akt antibodies. Compared with non-transgenic animals, phosphorylation of Akt on its activation sites Thr-308 and Ser-473 was significantly increased in cardioprotective PKC ϵ transgenic mice; b. Total Akt activity and PKC ϵ -associated Akt activity was measured by its ability to phosphorylate GSK-3 β . Total cardiac cellular Akt activity was increased in PKC ϵ cardioprotected mice when compared with non-transgenic mice. Moreover, PKC ϵ -associated Akt activity had greater elevation when compared with the increase in the total lysate pool.

a.



b.






Figure 23. Effect of PKC ϵ on the phosphorylation of eNOS *in vivo*. Non-transgenic and PKC ϵ transgenic mouse hearts were homogenized and separated with SDS-PAGE followed by immunoblotting with the eNOS and site-specific phospho-eNOS Ser¹¹⁷⁷ antibodies. Compared with non-transgenic animals, phosphorylation of eNOS on serine-1177, the Akt-specific activation site, was also enhanced in the PKC ϵ transgenic mice.



REFERENCES

1. Adams JA. Kinetic and catalytic mechanisms of protein kinases *Chem Rev* 101:2271-2290, 2001.
2. Akita Y. Protein kinase C- ϵ (PKC ϵ): its unique structure and function. *J Biochem.* 132: 847-852, 2002.
3. Alberts B. The cell as a collection of protein machines: preparing the next generation of molecular biologists. *Cell* 92:291-294, 1998.
4. Alderton WK, Cooper CE, Knowles RG. Nitric oxide synthases: structure, function and inhibition. *Biochem J* 357:593-615, 2001.
5. Alessi DR, James SR, Downes CP, Holmes AB, Gaffney PR, Reese CB, Cohen P. Characterization of a 3-phosphoinositide-dependent protein kinase which phosphorylates and activates protein kinase Balpha. *Curr Biol* 7:261-269, 1997.
6. Allen NP, Huang L, Burlingame A, Rexach M. Proteomic analysis of nucleoporin interacting proteins. *J Biol. Chem.* 276:29268-29274, 2001.
7. Altomare DA, Guo K, Cheng JQ, Sonoda G, Walsh K, Testa JR. Cloning, chromosomal localization and expression analysis of the mouse Akt2 oncogene. *Oncogene* 11:1055-1060, 1995.
8. Andjelkovic M, Alessi DR, Meier R, Fernandez A, Lamb NJ, Frech M, Cron P, Cohen P, Lucocq JM, Hemmings BA. Role of translocation in the activation and function of protein kinase B. *J Biol Chem* 272:31515-31524, 2001.
9. Andrew PJ, Mayer B. Enzymatic function of nitric oxide synthases. *Cardiovasc Res* 43: 521-531, 1999.
10. Armour KE, Armour KJ, Gallagher ME. Defective bone formation and anabolic response to exogenous estrogen in mice with targeted disruption of endothelial nitric oxide synthase. *Endocrinology* 142: 760-766, 2001.
11. Cohen MV, Baines CP, Downey JM. Ischemic preconditioning: from adenosine receptor of KATP channel. *Annu Rev Physiol.* 62:79-109, 2000.
12. Baines CP, Zhang J, Wang GW, Zheng YT, Xiu JX, Cardwell EM, Bolli R, Ping P. Mitochondrial PKCepsilon and MAPK form signaling modules in the murine heart: enhanced mitochondrial PKCepsilon-MAPK interactions and differential MAPK activation in PKCepsilon-induced cardioprotection. *Circ Res* .90:390-397, 2002.
13. Ballif BA, Shimamura A, Pae E, Blenis J. Disruption of 3-phosphoinositide-dependent kinase 1 (PDK1) signaling by the anti-tumorigenic and anti-proliferative agent n-alpha-tosyl-l-phenylalanyl chloromethyl ketone. *J Biol Chem* 276:12466-12475, 2001.

14. Balligand JL, Kobzik L, Han X, *et al.* Nitric oxide-dependent parasympathetic signaling is due to activation of constitutive endothelial (type III) nitric oxide synthase in cardiac myocytes. *J Biol Chem* 1995; 270: 14 852-14 856. *Biochem J* 299: 247–252, 1994.
15. Belevych AE, Harvey RD. Muscarinic inhibitory and stimulatory regulation of the L-type Ca^{2+} current is not altered in cardiac ventricular myocytes from mice lacking endothelial nitric oxide synthase. *J Physiol*. 528 :279-289, 2000.
16. Bell RM, Yellon DM. Bradykinin limits infarction when administered as an adjunct to reperfusion in mouse heart: the role of PI3K, Akt and eNOS. *J Mol Cell Cardiol* 35:185-193, 2003.
17. Bell AW, Ward MA, Blackstock WP, Freeman HN, Choudhary JS, Lewis AP, Chotai D, Fazel A, Gushue JN, Paiement J, Palcy S, Chevet E, Lafreniere-Roula, M., Solari, R., Thomas, D.Y., Rowley, A., Bergeron, J.J. Proteomics characterization of abundant Golgi membrane proteins. *J Biol Chem*. 276, 5152-516, 2001.
18. Bellacosa A, Chan TO, Ahmed NN, Datta K, Malstrom S, Stokoe D, McCormick F, Feng J, Tsichlis P. Akt activation by growth factors is a multiple-step process: the role of the PH domain. *Oncogene* 17:313-325, 1998.
19. Bellacosa A, Testa JR, Staal SP, Tsichlis PN. A retroviral oncogene, akt, encoding a serine-threonine kinase containing an SH2-like region. *Science* 11;254:274-277, 1999.
20. Benjamin IJ, McMillan DR. Stress (heat shock) proteins: molecular chaperones in cardiovascular biology and disease. *Circ Res* 83:117-132, 1998.
21. Bernardo NL, D'Angelo M, Okubo S, Joy A, Kukreja RC. Delayed ischemic preconditioning is mediated by opening of ATP-sensitive potassium channels in the rabbit heart. *Am J Physiol* 276:H1323-1330, 1999.
22. Berry DA, Keogh A, Dos Remedios CG. Nuclear membrane proteins in failing human dilated cardiomyopathy. *Proteomics*. 1:1507-1512, 2001.
23. Bolli R. The late phase of preconditioning. *Circ Res*. 87:972-983, 2002.
24. Brahmajothi MV, Campbell DL. Heterogeneous basal expression of nitric oxide synthase and superoxide dismutase isoforms in mammalian heart : implications for mechanisms governing indirect and direct nitric oxide-related effects. *Circ Res*. 85, 575-587, 1999.
25. Braunwald, Zipes, Libby. Heart disease: a textbook of cardiovascular medicine 6th edition. 2001.
26. Brazil DP, Hemmings BA. Ten years of protein kinase B signaling: a hard Akt to follow. *Trends Biochem Sci* 26:657-664, 2001.
27. Bredt DS, Hwang PM, Snyder SH. Localization of nitric oxide synthase indicating a neural role for nitric oxide. *Nature* 347: 768-770, 1990.
28. Brunet A, Datta SR, Greenberg ME. Transcription-dependent and -independent control of neuronal survival by the PI3K-Akt signaling pathway. *Curr Opin Neurobiol* 11:297-305, 2001.

29. Brunner F, Maier R, Andrew P, Wolkart G, Zechner R, Mayer B. Attenuation of myocardial ischemia/reperfusion injury in mice with myocyte-specific overexpression of endothelial nitric oxide synthase. *Cardiovasc Res* 57:55-62, 2003.
30. Busconi L, Michel T. Endothelial nitric oxide synthase: N-terminal myristoylation determines subcellular localization. *J Biol Chem* 268: 8410–8413, 1993.
31. Bustamante J, Bersier G, Badin RA, Cymeryng C, Parodi A, Boveris A. Sequential NO production by mitochondria and endoplasmic reticulum during induced apoptosis. *Nitric Oxide* 6: 333-341, 2002.
32. Castrillo A, Pennington DJ, Otto F, Parker PJ, Owen MJ, Bosca L. Protein kinase C epsilon is required for macrophage activation and defense against bacterial infection. *J. Exp. Med.* 194:1231–1242, 2001.
33. Cenni V, Doppler H, Sonnenburg ED, Maraldi N, Newwton AC, Toker A. Regulation of novel protein kinase C epsilon by phosphorylation. *Biochem. J.* 363:537-545, 2002.
34. Chan TO, Tsichlis PN. PDK2: A complex tail in one Akt. *Sci STKE.* 23;(66):PE1, 2001.
35. Chen R, Kim O, Yang J, Sato K, Eisenmann KM, McCarthy J, Chen H, Qiu Y. Regulation of Akt/PKB activation by tyrosine phosphorylation. *J Biol Chem* 276:31858-31862, 2001.
36. Chen J, Rappsilber J, Chiang YC, Russell P, Mann M, Denis CL. Purification and characterization of the 1.0 MDa CCR4-NOT complex identifies two novel components of the complex. *J Mol Biol* 314:683-694, 2001.
37. Chilov D, Camenisch G, Kvietikova I, Ziegler U, Gassmann M, Wenger RH. Induction and nuclear translocation of hypoxia-inducible factor-1 (HIF-1): heterodimerization with ARNT is not necessary for nuclear accumulation of HIF-1alpha. *J Cell Sci.* 112:1203-1212, 1999.
38. Coffey PJ, Woodgett JR. Molecular cloning and characterization of a novel putative protein-serine kinase related to the cAMP-dependent and protein kinase C families *Eur J Biochem* 201:475-481, 1991.
39. Condorelli G, Drusco A, Stassi G, Bellacosa A, Roncarati R, Iaccarino G, Russo MA, Gu Y, Dalton N, Chung C, Latronico MV, Napoli C, Sadoshima J, Croce CM, Ross J Jr. Akt induces enhanced myocardial contractility and cell size *in vivo* in transgenic mice. *Proc Natl Acad Sci U.S.A.* 99:12333-12338, 2002.
40. Cook SA, Matsui T, Li L, Rosenzweig A. Transcriptional effects of chronic Akt activation in the heart. *J Biol Chem.* 277:22528-22533, 2002.
41. Csukai M, Chen CH., De Matteis MA., Mochly-Rosen D. The coatomer protein beta'-COP, a selective binding protein (RACK) for protein kinase C epsilon. *J Biol Chem.* 272:29200-29206, 1997.
42. Datta SR, Brunet A, Greenberg ME. Cellular survival: a play in three Akts. *Genes Dev* 13:2905-2927, 1999.
43. Davda RK, Chandler LJ, and Guzman NJ. Protein kinase C modulates receptor-independent activation of endothelial nitric oxide synthase. *Eur J Pharmacol* 266:237–244, 1994.

44. Dimmeler S, Fleming I, Fisslthaler B, Hermann C, Busse R, Zeiher AM. Activation of nitric oxide synthase in endothelial cells by Akt-dependent phosphorylation. *Nature* 399:601-605, 1999.
45. Dinerman JL, Dawson TM, Schell MJ, Snowman A, Snyder SH. Endothelial nitric oxide synthase localized to hippocampal pyramidal cells: implications for synaptic plasticity. *Proc Natl Acad Sci U.S.A.* 91: 4214-4218, 1994.
46. Ding L, Wang H, Lang W, Xiao L. Protein kinase C-epsilon promotes survival of lung cancer cells by suppressing apoptosis through dysregulation of the mitochondrial caspase pathway. *J Biol Chem* 277:35305-35313, 2002.
47. Doble BW, Ping P, Kardami E. The epsilon subtype of protein kinase C is required for cardiomyocyte connexin-43 phosphorylation. *Circ Res.* 86, 293-301, 2000.
48. Doornbos RP, Theelen M, van der Hoeven PC, van Blitterswijk WJ, Verkleij AJ, van Bergen en Henegouwen PM. Protein kinase C zeta is a negative regulator of protein kinase B activity. *J Biol Chem* 274:8589-8596, 1999.
49. Dorn 2nd GW, Souroujon MC, Liron T, Chen C-H, Gray MO, Zhou HZ, Csukai M, Wu G, Lorenz JN, Mochly-Rosen D. Sustained in vivo cardiac protection by a rationally designed peptide that causes ϵ protein kinase C translocation. *Proc Natl Acad Sci USA* 96:12798-12803, 1999.
50. Dorn G.W., 2nd, Mochly-Rosen, D. Intracellular transport mechanisms of signal transducers. *Annu. Rev. Physiol.* 64: 407–429, 2002.
51. Du XL, Edelstein D, Dimmeler S, Ju Q, Sui C, Brownlee M. Hyperglycemia inhibits endothelial nitric oxide synthase activity by posttranslational modification at the Akt site. *J Clin Invest* 108:1341-1348, 2002.
52. Edmondson RD, Vondriska TM, Biedermant K, Zhang J, Jones R, Zheng YT, Allen DL, Xiu JX, Cardwell EM, Pisano MR, Ping P. Protein kinase C epsilon signaling complexes include metabolism- and transcription/translation-related proteins: complimentary separation techniques with LC/MS/MS. *Molecular Cellular Proteomics.* 6:421-433, 2002.
53. Faller DV. Endothelial cell responses to hypoxic stress. *Clin Exp Pharmacol Physiol.* 26:74-84, 1999.
54. Feng Y, Venema VJ, Venema RC, Tsai N, Caldwell RB. VEGF induces nuclear translocation of Flk-1/KDR, endothelial nitric oxide synthase, and caveolin-1 in vascular endothelial cells *Biochem Biophys Res Commun* 256:192-197, 1999.
55. Fisslthaler B, Dimmeler S, Hermann C, Busse R, and Fleming I. Phosphorylation and activation of the endothelial nitric oxide synthase by fluid shear stress. *Acta Physiol Scand.* 168: 81-88, 2000.
56. Fleming I, Busse R. Signal transduction of eNOS activation. *Cardiovasc Res.* 15;43:532-541, 1999.
57. Fleming I, Bauersachs J, Fisslthaler B, and Busse R. Ca^{2+} -independent activation of the endothelial nitric oxide synthase in response to tyrosine phosphatase inhibitors and fluid shear stress. *Circ Res.* 82: 686–695, 1998.

58. Fleming I, Fisslthaler B, Dimmeler S, Kemp BE, and Busse R. Phosphorylation of Thr495 regulate Ca^{2+} /calmodulin- dependent endothelial nitric oxide synthase activity. *Circ Res* 88: 68–75, 2001
59. Flynn P, Wongdagger M, Zavar M, Dean NM, Stokoe D. Inhibition of PDK-1 activity causes a reduction in cell proliferation and survival. *Curr Biol* 10:1439-1442, 2002.
60. Fontana J, Fulton D, Chen Y, Fairchild TA, McCabe TJ, Fujita N, Tsuruo T, Sessa WC. Domain mapping studies reveal that the M domain of hsp90 serves as a molecular scaffold to regulate Akt-dependent phosphorylation of endothelial nitric oxide synthase and NO release. *Circ Res* 90:866-873, 2002.
61. Fraser ID, Cong M, Kim J, Rollins EN, Daaka Y, Lefkowitz RJ, Scott JD. Assembly of an A-kinase-anchoring protein- β_2 -adrenergic receptor complex facilitates receptor phosphorylation and signaling. *Curr Biol* 10:409-412, 2001.
62. Fujio Y, Nguyen T, Wencker D, Kitsis RN, Walsh K. Akt promotes survival of cardiomyocytes in vitro and protects against ischemia-reperfusion injury in mouse heart *Circulation* 101:660-667, 2001.
63. Fulton D, Gratton JP, McCabe TJ, Fontana J, Fujio Y, Walsh K, Franke TF, Papapetropoulos A, Sessa WC. Regulation of endothelium-derived nitric oxide production by the protein kinase Akt. *Nature* 399:597-601, 1999.
64. Gallis B, Corthals GL, Goodlett DR, Ueba H, Kim F, Presnell SR, Figes D, Harrison DG, Berk BC, Aebersold R, and Corson MA. Identification of flow-dependent endothelial nitric oxide synthase phosphorylation sites by mass spectrometry and regulation of phosphorylation and nitric oxide production by the phosphatidylinositol 3-kinase inhibitor LY294002. *J Biol Chem* 274: 30101–30108, 1999.
65. Garcia-Cardena G, Fan R, Shah V, Sorrentino R, Cirino G, Papapetropoulos A, Sessa WC. Dynamic activation of endothelial nitric oxide synthase by Hsp90. *Nature* 392:821-824, 1998.
66. Gavin AC, Bosche M, Krause R, Grandi P, Marzioch M, Bauer A, Schultz J, Rick JM, Michon AM, Cruciat CM, Remor M, Hofert C, Schelder M, Brajenovic M, Ruffner H, Merino A, Klein K, Hudak M, Dickson D, Rudi T, Gnau V, Bauch A, Bastuck S, Huhse B, Leutwein C, Heurtier MA, Copley RR, Edelmann A, Querfurth E, Rybin V, Drewes G, Raida M, Bouwmeester T, Bork P, Seraphin B, Kuster B, Neubauer G, Superti-Furga G. Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature* 415:141-147, 2002.
67. Gerber HP, McMurtrey A, Kowalski J, Yan M, Keyt BA, Dixit V, Ferrara N. Vascular endothelial growth factor regulates endothelial cell survival through the phosphatidylinositol 3'-kinase/Akt signal transduction pathway. Requirement for Flk-1/KDR activation. *J Biol Chem* 273:30336-30343, 1998.
68. Gilliam MB, Sherman MP, Griscavage JM, Ignarro LJ. A spectrophotometric assay for nitrate using NADPH oxidation by *Aspergillus* nitrate reductase. *Anal Biochem.* 212:359-365, 1993.
69. Gingras AC, Gygi SP, Raught B, Polakiewicz RD, Abraham RT, Hoekstra MF, Aebersold R, Sonenberg N. Regulation of 4E-BP1 phosphorylation: a novel two-step mechanism. *Genes Dev* 13:1422-37, 1999.

70. Godecke A, Heinicke T, Kamkin A, Kiseleva I, Strasser RH, Decking UK, Stumpe T, Isenberg G, Schrader J. Godecke A, Heinicke T, Kamkin A, Kiseleva I, Strasser RH, Decking UK, Stumpe T, Isenberg G, Schrader J. Inotropic response to beta-adrenergic receptor stimulation and anti-adrenergic effect of ACh in endothelial NO synthase-deficient mouse hearts. *J Physiol*. 532:195-204, 2001.
71. Godovac-Zimmermann J, Soskic V, Poznanovic S, Brianza F. Functional proteomics of signal transduction by membrane receptors. *Electrophoresis* 20: 952-961, 1999.
72. Graness A, Adomeit A, Heinze R, Wetzker R, Liebmann C. A novel mitogenic signaling pathway of bradykinin in the human colon carcinoma cell line SW-480 involves sequential activation of a Gq/11 protein, phosphatidylinositol 3-kinase beta, and protein kinase C epsilon. *J. Biol. Chem*. 273:32016–32022, 1998.
73. Graves PR, Haystead TA. A functional proteomics approach to signal transduction. *Recent Prog Horm Res*. 58:1-24, 2003.
74. Gravin AC. Functional organization of the yeast proteome by systematic analysis of protein complexes. *Nature*. 415:141-147, 2002.
75. Gray MO, Karliner JS, Mochly-Rosen D. A selective ϵ -protein kinase C antagonist inhibits protection of cardiac myocytes from hypoxia-induced cell death. *J Biol Chem* 272: 30945-30953, 1997.
76. Gross SS, Wolin MS. Nitric oxide: pathophysiological mechanisms. *Annu Rev Physiol* 57: 737-769, 1995.
77. Gross A, Jockel J, Wei MC, Korsmeyer SJ. Enforced dimerization of BAX results in its translocation, mitochondrial dysfunction and apoptosis. *EMBO J* 171:3878-3885, 1998.
78. Guo Y, Jones WK, Xua YT, Tang XL, Bao W, Wu WJ, Han H, Laubach VE, Ping P, Yang Z, Qiu Y, Bolli R. The late phase of ischemic preconditioning is abrogated by targeted disruption of the inducible NO synthase gene. *Proc Natl Acad Sci U S A* 96:11507-11512, 1999.
79. Gyurko R, Kuhlencordt P, Fishman MC, Huang PL. Modulation of mouse cardiac function *in vivo* by eNOS and ANP. *Am J Physiol* 278:H971-981, 2000.
80. Hare JM, Lofthouse RA, Juang GJ, Colman L, Ricker KM, Kim B, Senzaki H, Cao S, Tunin RS, Kass DA. Contribution of caveolin protein abundance to augmented nitric oxide signaling in conscious dogs with pacing-induced heart failure. *Circ Res*. 86:1085-1092, 2000.
81. Harrison PM, Kumar A, Lang N, Snyder M, Gerstein M. A question of size: the eukaryotic proteome and the problem in defining it. *Nucleic Acids Res*. 30: 1083-1090, 2002.
82. Hartwell LH, Hopfield JJ, Leibler S, Murry AW. From molecular to modular cell biology *Nature* (Supp). 402:C47-C52, 1999.
83. Hemmens B, Mayer B. Enzymology of nitric oxide synthases. *Methods Mol Biol* 100:1-32, 1998.
84. Hill MM, Feng J, Hemmings BA. Identification of a plasma membrane Raft-associated PKB Ser473 kinase activity that is distinct from ILK and PDK1. *Curr Biol* 12:1251-1255, 2002.

85. Hirata K, Kuroda R, Sakoda T, Katayama M, Inoue N, Suematsu M, Kawashima S, and Yokoyama M. Inhibition of endothelial nitric oxide synthase activity by protein kinase C. *Hypertension*. 25:180-185, 1995.
86. Ho Y, Ho Y, Gruhler A, Heilbut A, Bader GD, Moore L, Adams SL, Millar A, Taylor P, Bennett K, Boutilier K, Yang L, Wolting C, Donaldson I, Schandorff S, Shewnarane J, Vo M, Taggart J, Goudreau M, Muskat B, Alfaro C, Dewar D, Lin Z, Michalickova K, Willems AR, Sassi H, Nielsen PA, Rasmussen KJ, Andersen JR, Johansen LE, Hansen LH, Jespersen H, Podtelejnikov A, Nielsen E, Crawford J, Poulsen V, Sorensen BD, Matthies J, Hendrickson RC, Gleeson F, Pawson T, Moran MF, Durocher D, Mann M, Hogue CW, Figeys D, Tyers M. Systematic identification of protein complexes in *Saccharomyces cerevisiae* by mass spectrometry. *Nature* 415:180-183, 2002.
87. Hoang VM, Foulk R, Clauser K, Burlingame A, Gibson BW, Fisher SJ. Functional proteomics: examining the effects of hypoxia on the cytotrophoblast protein repertoire. *Biochemistry* 40:4077-4086, 2001.
88. Hodgkinson CP, Sale EM, Sale GJ. Characterization of PDK2 activity against protein kinase B gamma. *Biochemistry* 41:10351-10359, 2002.
89. Hoffman A, Gloe T, Pohl U. Hypoxia-induced upregulation of eNOS gene expression is redox-sensitive: a comparison between hypoxia and inhibitors of cell metabolism. *J Cell Physiol*. 188: 33-44, 1998.
90. Hoffmann P, Ji H, Moritz RL, Connolly LM, Frecklington DF, Layton MJ, Eddes JS, Simpson RJ. Continuous free-flow electrophoresis separation of cytosolic proteins from the human colon carcinoma cell line LIM 1215: a non two-dimensional gel electrophoresis-based proteome analysis strategy. *Proteomics* 1:807-818, 2001.
91. Hondermarck H, Vercoutter-Edouart AS, Revillion F, Lemoine J, El-Yazidi-Belkoura I, Nurcombe V, Peyrat JP. Proteomics of breast cancer for marker discovery and signal pathway profiling. *Proteomics*. 1:1216-1232, 2001.
92. Huang PL. Mouse models of nitric oxide synthase deficiency. *J Am Soc Nephrol* 16:S120-123, 2000.
93. Huang C, Zhou Q, Liang P, Hollander MS., Sheikh F, Li X, Greaser M, Shelton GD, Evans S, Chen J. Characterization and *in vivo* functional analysis of splice variants of Cypher. *J Biol Chem*. 278:7360-7365, 2003.
94. Hubbard MJ, Cohen P. On target with a new mechanism for the regulation of protein phosphorylation. *Trends Biochem Sci* 18:172-177, 1993.
95. Huber LA, Pfaller K, Vietor I. Organelle Proteomics Implication for subcellular fraction in proteomics. *Circ. Res.* 92: 962-968, 2003.
96. Huber LA. Opinion: is proteomics heading in the wrong direction? *Nat Rev Mol Cell Biol*. 4: 74-80, 2003.
97. Hunter T. Signaling--2000 and beyond. *Cell* 100: 113-127, 2000.
98. Husi H, Grant SG. Isolation of 200-kDa complexes of NMDA receptor and postsynaptic density 95 from mouse brain. *J Neurochem* 277:281-291, 2001.
99. Husi H, Ward MA, Choudhary JS, Blackstock WP, Grant SG. Proteomic analysis of NMDA receptor-adhesion protein signaling complexes. *Nat Neurosci*. 3:661-669, 2000.

100. Jones PF, Jakubowicz T, Pitossi FJ, Maurer F, Hemmings BA. Molecular cloning and identification of a serine/threonine protein kinase of the second-messenger subfamily. *Proc Natl Acad Sci U.S.A.* 88:4171-4175, 1991.
101. Jordan JD, Landau EM, Iyengar R. Signaling networks: the origins of cellular multitasking. *Cell* 103:193-200, 2000.
102. Ju H, Venema VJ, Marrero MB, Venema RC. Inhibitory interactions of the bradykinin B2 receptor with endothelial nitric-oxide synthase. *J Biol Chem.* 273:24025-24029, 1998.
103. Kandel ES, Hay N. The regulation and activities of the multifunctional serine/threonine kinase Akt/PKB. *Exp Cell Res* 253:210-29, 1999.
104. Kane LP, Shapiro VS, Stokoe D, Weiss A. Induction of NF-kappaB by the Akt/PKB kinase. *Curr Biol* 9:601-4, 1999.
105. Kelly RA, Balligand JL, Smith TW. Nitric oxide and cardiac function. *Circ Res* 79:363-80, 1996.
106. Khasar SG, Lin YH, Martin A, Dadgar J, McMahon T, Wang D, Hundle B, Aley KO, Isenberg W, McCarter G, Green PG, Hodge CW, Levine JD, Messing RO. A novel nociceptor signaling pathway revealed in protein kinase C epsilon mutant mice. *Neuron* 24:253-260, 1999.
107. Kim F, Gallis B, and Corson MA. TNF inhibits flow and insulin signaling leading to NO production in aortic endothelial cells. *Am J Physiol Cell Physiol* 280: C1057-C1065, 2001.
108. Klatt P, Pfeiffer S, List BM, *et al.* Characterization of the heme-deficient neuronal nitric oxide synthase reveals a role for heme in subunit dimerization and binding of the amino acid substrate and tetrahydrobiopterin. *J Biol Chem.* 271: 7336-7342, 1996.
109. Knezevic V, Leethanakul C, Bichsel VE, Worth JM, Prabhu VV, Gutkind JS, Liotta LA, Munson PJ, Petricoin EF. 3rd, Krizman DB. Proteomic profiling of the cancer microenvironment by antibody arrays. *Proteomics* 1:1271-1278, 2001.
110. Konoshi H, Kuroda S, Tanaka M, Matsuzaki H, Ono Y, Kameyama K, Haga T, Kikkawa U. Molecular cloning and characterization of a new member of the RAC protein kinase family: Association of the pleckstrin homology domain of three types of RAC protein kinase with protein kinase C subspecies and beta gamma subunits of G proteins. *Biochem Biophys. Res. Commun.* 216: 526-534, 1995.
111. Lantinho RL, Rosenfeld CR, Yuhanna IS, German Z, Chen Z, and Shaul PW. Estrogen acutely stimulates nitric oxide synthase activity in fetal pulmonary artery endothelium. *Am J. Physiol.* 274:R797-807, 1998.
112. Lehel C, Olah Z, Jakab G, Szallasi Z, Petrovics G, Harta G, Blumberg PM, Anderson WB. Protein kinase C epsilon subcellular localization domains and proteolytic degradation sites. A model for protein kinase C conformational changes. *J. Biol. Chem.* 270: 19651-19658, 1995.
113. Lewis TS, Hunt JB, Aveline LD, Jonscher KR, Louie DF, Yeh JM, Nahreini TS, Resing KA, Ahn NG. Identification of novel MAP kinase pathway signaling targets by functional proteomics and mass spectrometry. *Mol Cell* 6:1343-1354, 2000.

114. Li W, Zhang J, Flechner L, Hyun T, Yam A, Franke TF, Pierce JH. Protein kinase C- α overexpression stimulates Akt activity and suppresses apoptosis induced by interleukin 3 withdrawal. *Oncogene* 18:6564-6572, 1999.
115. Li RC, Ping P, Zhang J, Wead WB, Cao X, Gao J, Zheng Y, Huang S, Han J, Bolli R. PKC ϵ modulates NF- κ B and AP-1 via mitogen-activated protein kinases in adult rabbit cardiomyocytes. *Am J Physiol*. 279:H1679-1689, 2000.
116. List BM, Klösch B, Volker C. Characterization of bovine endothelial nitric oxide synthase as a homodimer with down-regulated uncoupled NADPH oxidase activity: tetrahydrobiopterin binding kinetics and role of haem in dimerization. *Biochem J* 323:159-165, 1997.
117. Liu GS, Cohen MV, Mochly-Rosen D, Downey JM. Protein kinase C ϵ is responsible for the protection of preconditioning in rabbit cardiomyocytes. *J Mol Cell Cardiol* 31:1937-1948, 1999.
118. Lopez MF, Melov S. Applied proteomics: mitochondrial proteins and effect on function. *Circ Res*. 90:380-389, 2002.
119. Mao M, Fang X, Lu Y, Lapushin R, Bast RC Jr, Mills GB. Inhibition of growth-factor-induced phosphorylation and activation of protein kinase B/Akt by atypical protein kinase C in breast cancer cells. *Biochem J*. 352:475-478, 2002.
120. Matsubara M, Titani K, and Taniguchi H. Interaction of calmodulin-binding domain peptides of nitric oxide synthase with membrane phospholipids: regulation by protein phosphorylation and Ca²⁺-calmodulin. *Biochemistry* 35: 14651–14658, 1996.
121. Matsui T, Tao J, del Monte F, Lee KH, Li L, Picard M, Force TL, Franke TF, Hajjar RJ, Rosenzweig A. Akt activation preserves cardiac function and prevents injury after transient cardiac ischemia *in vivo*. *Circulation* 104:330-335, 2001.
122. McQuillan LP, Leung GK, Marsden PA, Kostyk SK, and Kourembanas S. Hypoxia inhibits expression of eNOS via transcriptional and posttranscriptional mechanisms. *Am J Physiol* 267: H1921–1927, 1994.
123. Michell BJ, Chen ZP, Tiganis T, Stapleton D, Katsis F, Power DA, Sim AT, Kemp BE. Coordinated control of endothelial nitric-oxide synthase phosphorylation by protein kinase C and the cAMP-dependent protein kinase. *J Biol Chem* 276:17625-17628, 2001.
124. Michell BJ, Griffiths JE, Mitchelhill KI, Rodriguez-Crespo I, Tiganis T, Bozinovski S, de Montellano PR, Kemp BE, Pearson RB. The Akt kinase signals directly to endothelial nitric oxide synthase. *Curr Biol*. 9:845-848, 1999.
125. Mohamed F, Monge JC, Gordon A, Cernacek P, Blais D, and Stewart DJ. Lack of role for nitric oxide (NO) in the selective destabilization of endothelial NO synthase mRNA by tumor necrosis factor- α . *Arterioscler Thromb Vasc Biol* 15: 52–57, 1995.
126. Molloy MP, Herbert BR, Slade MB, Rabilloud T, Nouwens AS, Williams KL., Gooley AA. Proteomic analysis of the Escherichia coli outer membrane *Eur J Biochem* 267: 871-2881, 2001.
127. Moriya S, Kazlauskas A, Akimoto K, Hirai S, Mizuno K, Takenawa T, Fukui Y, Watanabe Y, Ozaki S, Ohno S. Platelet-derived growth factor activates protein kinase C ϵ through redundant and independent signaling pathways involving phospholipase C γ or phosphatidylinositol 3-kinase. *Proc. Natl. Acad. Sci. USA* 93:151–155, 1999.

128. Murry CE, Jennings RB, Reimer KA. Preconditioning with ischemia: a delay of lethal cell injury in ischemic myocardium. *Circulation*, 74:1124-1136, 1986.
129. Newton AC. Regulation of the ABC kinases by phosphorylation: protein kinase C as a paradigm. *Biochem. J.* 370:361-371, 2003.
130. O'Rourke B. Myocardial K_{ATP} channels in preconditioning. *Circ Res.* 87:845-855, 2000.
131. Oh H, Fujio Y, Kunisada K, Hirota H, Matsui H, Kishimoto T, Yamauchi-Takahara K. Activation of phosphatidylinositol 3-kinase through glycoprotein 130 induces protein kinase B and p70 S6 kinase phosphorylation in cardiac myocytes. *J Biol Chem* 273:9703-9710, 1998.
132. Ohi MD, Link AJ, Ren L, Jennings JL., McDonald WH., Gould KL. Proteomics analysis reveals stable multiprotein complexes in both fission and budding yeasts containing Myb-related Cdc5p/Cef1p, novel pre-mRNA splicing factors, and snRNAs. *Mol Cell Biol.* 22:2011-2024, 2002.
133. Palzkill T. Proteomics. Kluwer Academic Publishers. P1-4. 2001.
134. Pandey A, Mann M. Proteomic to study genes and genomes. *Nature* 405:837-846, 2000.
135. Pass JM, Zhang J, Vondriska TM, Ping P. Functional proteomic analysis of the protein kinase C signaling system. *Methods Mol Biol.* 233:369-385, 2003.
136. Pass JM, Zheng Y, Wead WB, Zhang J, Li RC, Bolli R Ping P. PKCepsilon activation induces dichotomous cardiac phenotypes and modulates PKCepsilon-RACK interactions and RACK expression. *Am J Physiol.* 88:H946-955, 2001.
137. Passier R, Richardson JA, Olson EN. Oracle, a novel PDZ-LIM domain protein expressed in heart and skeletal muscle *Mech Dev.* 92:277-284, 2000.
138. Persad S, Attwell S, Gray V, Mawji N, Deng JT, Leung D, Yan J, Sanghera J, Walsh MP, Dedhar S. Regulation of protein kinase B/Akt-serine 473 phosphorylation by integrin-linked kinase: critical roles for kinase activity and amino acids arginine 211 and serine 343. *J Biol Chem.* 276:27462-27469, 2001.
139. Phadke ND, Molloy MP, Steinhoff SA, Ulintz PJ, Andrews PC, Maddock JR. Analysis of the outer membrane proteome of *Caulobacter crescentus* by two-dimensional electrophoresis and mass spectrometry. *Proteomics* 1:705-720, 2001.
140. Ping P, Takano H, Zhang J, Tang XL, Qiu Y, Li RC-X, Banerjee S, Dawn B, Balafanova Z, Bolli R. Isoform-selective activation of protein kinase C by nitric oxide in the heart of conscious rabbits: a signaling mechanism for both nitric oxide-induced and ischemia-induced preconditioning. *Circ Res.* 84:587-604, 1999.
141. Ping P, Zhang J, Huang S, Cao X, Tang XL, Li RCX, Zheng YT, Qiu Y, Clerk A, Sugden PH, and Bolli R. PKC-dependent activation of p46/p54 JNKs during ischemic preconditioning in conscious rabbits. *Am J Physiol* 277:H1771-1785, 1999.
142. Ping P, Zhang J, Pierce WM Jr, Bolli R. Functional proteomic analysis of PKCε signaling complexes in the normal heart and during cardioprotection. *Circ Res* 88:59-62, 2001.

143. Ping P, Zhang J, Qiu Y, Tang XL, Manchikalapudi S, Cao X, Bolli R. Ischemic preconditioning induces selective translocation of PKC isoforms ϵ and η in the heart of conscious rabbits without subcellular redistribution of total PKC activity. *Circ Res* 81:404-414, 1997.
144. Ping P, Zhang J, Zheng Y, Li RCX, Dawn B, Tang XL, Takano H, Balafanova Z, Bolli R. Demonstration of selective protein kinase C-dependent activation of Src and Lck tyrosine kinases during ischemic preconditioning in conscious rabbits. *Circ Res* 85: 542-550, 1999.
145. Ping P, Song C, Zhang J, Guo Y, Cao X, Li RC, Wu W, Vondriska TM, Pass JM, Tang XL, Pierce WM, Bolli R. Formation of protein kinase C(epsilon)-Lck signaling modules confers cardioprotection. *J. Clin. Invest.* 109, 499–507, 2002.
146. Qin Q, Downey JM, Cohen MV. Acetylcholine but not adenosine triggers preconditioning through PI3-kinase and a tyrosine kinase *Am J Physiol.* 284:H727-734, 2003.
147. Qiu Y, Ping P, Tang XL, Manchikalapudi S, Rizvi A, Zhang J, Takano H, Wu WJ, Teschner S, Bolli R. Direct evidence that protein kinase C plays an essential role in the development of late preconditioning against myocardial stunning in conscious rabbits and that ϵ is the isoform involved. *J Clin Invest* 101:2182-2198, 1999.
148. Radomski MW, Palmer RM, Moncada S. An L-arginine/nitric oxide pathway present in human platelets regulates aggregation. *Proc Natl Acad Sci U. S. A.* 57: 5193-6197, 1990.
149. Ragolia L, Palaia T, Paric E, Maesaka JK. Elevated L-PGDS activity contributes to PMA-induced apoptosis concomitant with downregulation of PI3-K. *Am J Physiol* 284:C119-26, 2003.
150. Raman CS, Li H, Martasek P, Kral V, Masters BS, Poulos TL. Crystal structure of constitutive endothelial nitric oxide synthase: a paradigm for protein function involving a novel metal center. *Cell* 95: 939-950, 1998.
151. Reiner M, Bloch W, Addicks K. Functional interaction of caveolin-1 and eNOS in myocardial capillary endothelium revealed by immunoelectron microscopy. *J Histochem.* 49:1605-1610, 2001.
152. Sakoda T, Hirata K, Kuroda R, Miki N, Suematsu M, Kawashima S, and Yokoyama M. Myristoylation of endothelial cell nitric oxide synthase is important for extracellular release of nitric oxide. *Mol Cell Biochem* 152: 143–148, 1995.
153. Schagger H, Cramer WA, Jagow G. Analysis of molecular masses and oligomeric states of protein complexes by blue native electrophoresis and isolation of membrane protein complexes by two-dimensional native electrophoresis. *Analytical biochemistry.* 217: 220-230, 1994.
154. Schagger H. Quantification of oxidative phosphorylation enzymes after blue native electrophoresis and two-dimensional resolution: normal complex I protein amounts in Parkinson's disease conflict with reduced catalytic activities. *Electrophoresis.* 16:763-770, 1995.
155. Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell* 103:211-225, 2000.
156. Scott-Burden T, Engler DA, Tock CL, Schwarz JJ, Casscells SW. Liposomal induction of NO synthase expression in cultured vascular smooth muscle cells. *Biochem Biophys Res Commun.* 231:780-783, 1997.

157. Sessa WC, Barber CM, and Lynch KR. Mutation of *N*-myristoylation site converts endothelial cell nitric oxide synthase from a membrane to a cytosolic protein. *Circ Res* 72: 921–924, 1993.
158. Shinmura K, Tang XL., Wang Y, Xuan YT, Liu SQ, Takano H, Bhatnagar A, Bolli R. Cyclooxygenase-2 mediates the cardioprotective effects of the late phase of ischemic preconditioning in conscious rabbits. *Proc Natl Acad Sci U S A.* 97:10197-10202, 2000.
159. Smart EJ, Ying YS, Mineo C, Anderson RG. A detergent-free method for purifying caveolae membrane from tissue culture cells. *Proc Natl Acad Sci U S A.* 92:10104-10108, 1995.
160. Song C, Vondriska TM, Wang G, Klein JB, Cao X, Zhang J, Kang YJ, D'Souza S, Ping P. Molecular conformation dictates signaling module formation: example of PKC ϵ and Src tyrosine kinase. *Am J Physiol.* 282:H1166-1171, 2002.
161. Sonnenburg ED, Gao T, Newton AC. The phosphoinositide dependent kinase, PDK-1, phosphorylates conventional protein kinase C isozymes by a mechanism that is independent of phosphoinositide-3-kinase. *J. Biol. Chem.*, 276:45289-45297, 2001.
162. Stephens L, Anderson K, Stokoe D, Erdjument-Bromage H, Painter GF, Holmes AB, Gaffney PR, Reese CB, McCormick F, Tempst P, Coadwell J, Hawkins PT. Protein kinase B kinases that mediate phosphatidylinositol 3,4,5-trisphosphate-dependent activation of protein kinase B. *Science* 279:710-714, 1998.
163. Stokoe D, Stephens LR, Copeland T, Gaffney PR, Reese CB, Painter GF, Holmes AB, McCormick F, Hawkins PT. Dual role of phosphatidylinositol-3,4,5-trisphosphate in the activation of protein kinase B. *Science* 277:567-570, 1997.
164. Takahashi M, Mukai H, Oishi K, Isagawa T, Ono Y. Association of immature hypophosphorylated protein kinase c epsilon with an anchoring protein CG-NAP. *J. Biol. Chem.* 275:34592–34596, 2000.
165. Toker A, Newton AC. Akt/protein kinase B is regulated by autophosphorylation at the hypothetical PDK-2 site. *J Biol Chem* 275:8271-8274, 2000.
166. Tong H, Chen W, Steenbergen C, Murphy E. Ischemic preconditioning activates phosphatidylinositol-3-kinase upstream of protein kinase C. *Circ. Res.* 87:309–315, 2000.
167. Uemura K, Aki T, Yamaguchi K, Yoshida K. Protein kinase C-epsilon protects PC12 cells against methamphetamine-induced death: possible involvement of suppression of glutamate receptor. *Life Sci.* 21;72:1595-1607, 2003.
168. Van Eyk, J.E. Proteomics: unraveling the complexity of heart disease and striving to change cardiology. *Curr Opin Mol Ther.* 3: 546-553, 2001.
169. Venema RC, Ju H, Zou R, Ryan JW, Venema VJ. Subunit interactions of endothelial nitric-oxide synthase. Comparisons to the neuronal and inducible nitric-oxide synthase isoforms. *J Biol Chem* 272: 1276-1282, 1997.
170. Vondriska TM, Ping P. Functional proteomics to study protection of the ischaemic myocardium. *Expert Opin Ther Targets.* 6:563-570, 2002.

171. Vondriska TM, Zhang J, Ping P. Proteomic characterization of protein kinase C signaling tasks in the heart. *Proteomic and genomic analysis of cardiovascular disease*. Edited by Van Eyk JE, Dunn MJ. Wiley-VCH, p293-306, 2002.
172. Vondriska TM, Zhang J, Song C, Tang XL, Cao X, Baines CP, Pass JM, Wang S, Bolli R, Ping P. PKC ϵ -Src modules directs signal transduction in nitric oxide-induced cardioprotection: complex formation as a means for cardioprotective signaling. *Circ Res* 88:1306-1313, 2001.
173. Vondriska TM, Klein JB., Ping P. Use of functional proteomics to investigate PKC epsilon-mediated cardioprotection: the signaling module hypothesis. *Am J Physiol*. 280:H1434-1441, 2001.
174. Wallenstein S, Zucker CL, Fleiss JL. Some statistical methods used in circulation research. *Circ Res*. 47:1-9, 1980.
175. Wasinger VC, Cordwell SJ, Cerpa-Poljak A, Yan JX, Gooley AA, Wilkins MR, Duncan MW, Harris R, Williams KL, Humphery-Smith I. Progress with gene-product mapping of the Mollicutes: *Mycoplasma genitalium*. *Electrophoresis*. 16:1090-1094, 1995.
176. Wen HC, Huang WC, Ali A, Woodgett JR, Lin WW. Negative regulation of phosphatidylinositol 3-kinase and Akt signaling pathway by PKC. *Cell Signal*. 15:37-45, 2003.
177. Williams MR, Arthur JS, Balendran A, van der Kaay J, Poli V, Cohen P, Alessi DR. The role of 3-phosphoinositide-dependent protein kinase 1 in activating AGC kinases defined in embryonic stem cells. *Curr Biol* 10:439-448, 2002.
178. Wu CC, Taylor RS, Lane DR, Ladinsky MS, Weisz JA, Howell KE. GMx33: a novel family of trans-Golgi proteins identified by proteomics. *Traffic*, 1: 963-75, 2000.
179. Xi L, Kukreja RC. Pivotal role of nitric oxide in delayed pharmacological preconditioning against myocardial infarction. *Toxicology* 155:37-44, 2000.
180. Xu KY, Huso DL, Dawson TM, Bredt DS, Becker LC. Nitric oxide synthase in cardiac sarcoplasmic reticulum. *Proc Natl Acad Sci*. 96:657-662, 1999.
181. Xuan YT, Tang XL, Qiu Y, Banerjee S, Takano H, Han H, Bolli R. Biphasic response of cardiac NO synthase isoforms to ischemic preconditioning in conscious rabbits. *Am J Physiol*. 279:H2360-2371, 2000.
182. Yamashita K, Kajstura J, Discher DJ, Wasserlauf BJ, Bishopric NH, Anversa P, Webster KA. Reperfusion-activated Akt kinase prevents apoptosis in transgenic mouse hearts overexpressing insulin-like growth factor-1. *Circ Res*. 88:609-614, 2001.
183. Yang J, Corn P, Thompson V, Good VM, Hess D, Hemmings BA, Barford D. Molecular mechanism for the regulation of protein kinase B/Akt by hydrophobic motif phosphorylation. *Mol Cell* 9:1227-40, 2002.
184. Yet SF, Tian R, Layne MD, Wang ZY, Maemura K, Solovyeva M, Ith B, Melo LG, Zhang L, Ingwall JS, Dzau VJ, Lee ME, Perrella MA. Cardiac-specific expression of heme oxygenase-1 protects against ischemia and reperfusion injury in transgenic mice. *Circ Res*. 89:168-173, 2001.

185. Yoshizumi M, Perrella MA, Burnett JC Jr, and Lee ME. Tumor necrosis factor downregulates an endothelial nitric oxide synthase mRNA by shortening its half-life. *Circ Res* 73: 205–209, 1993.
186. Ytrehus K, Liu Y, Downey JM. Preconditioning protects ischemic rabbit heart by protein kinase C activation. *Am J Physiol* 266: H1145-1152, 1994.
187. Zhang J, Patel JM, Li YD, and Block ER. Proinflammatory cytokines downregulate gene expression and activity of constitutive nitric oxide synthase in porcine pulmonary artery endothelial cells. *Res Commun Mol Pathol Pharmacol* 96: 71–87, 1997.
188. Zhou HZ, Karliner JS, Gray MO. Moderate alcohol consumption induces sustained cardiac protection by activating PKC-epsilon and Akt. *Am J Physiol*. 283:H165-174, 2002.
189. Zhou Q, Chu PH, Huang C, Cheng CF, Martone ME, Knoll G, Shelton GD, Evans S, Chen J. Ablation of Cypher, a PDZ-LIM domain Z-line protein, causes a severe form of congenital myopathy. *J Cell Biol*. 155:605-612, 2001.
190. Ziegler WH, Parekh DB, Le Good JA, Whelan RD, Kelly JJ, Frech M, Hemmings BA, Parker PJ. Rapamycin-sensitive phosphorylation of PKC on a carboxy-terminal site by an atypical PKC complex. *Curr Biol* 9:522-529, 1999.

CURRICULUM VITAE

Name: Jun Zhang

Present Address: Department of Physiology & Biophysics,
Health science center, A Bldg., RM 1105
University of Louisville, KY, 40202

Phone: (502) 339-9232 (Home)
(502) 852-5379 (Office)

Fax: (502) 852-6239

Email: Julyjun@yahoo.com

EDUCATION

08/2001-Present: Doctoral student, Department of Physiology/Biophysics,
University of Louisville. U.S.A. Major: Physiology

08/1991 - 07/1994: Master of Science, Institute of Microbiology,
Chinese Academy of Sciences. P.R.China. Major: Biochemistry.

09/1987 - 07/1991: Bachelor of Science, Dept. of Microbiology,
Shandong University, P.R.China. Major: Microbiology.

EMPLOYMENT AND WORKING EXPERIENCE

01/1996 – 12/2002: Senior Research Associate
Dept. of Physiology and Biophysics / Division of Cardiology,
School of Medicine, University of Louisville. U.S.A.

08/1994 - 12/1995: Research Associate
Dept. of Molecular Biology, Institute of Microbiology,
Chinese Academy of Sciences. P.R.China.

HONORS AND AWARDS

1. 2001: Trainee Abstract Award: American Heart Association, Scientific Sessions, Anaheim, CA.

2. 1993: Third Prize in Young Scientific Research Presentation of Chinese Academy of Sciences, P.R.China.

SOCIETY MEMBERSHIP

Member, International Society for Heart Research.

Fellow, American Heart Association, Basic Sciences Council

PUBLICATIONS

1. **Zhang J**, Baines CP, Vondriska TM, Cardwell EM, Zong C, Wang GW, Ping P. Characterization of myocardial PKC ϵ -Akt-eNOS signaling modules via functional proteomic analysis. (in preparation)
2. **Zhang J**, Baines CP, Vondriska TM, Zong C, Cardwell EM, Wang GW, Loo R, Ping P. Functional proteomic mapping of the myocardial PKC ϵ signaling subproteome in the mouse heart: a subcellular compartment study. (Submitted to *Proteomics*)
3. **Zhang J**, Ping P, Vondriska TM, Tang XL, Wang GW, Cardwell EM, Bolli R. Cardioprotection involves activation of NF- κ B via PKC-dependent tyrosine and serine phosphorylation of I κ B- α . *Am J Physiol*. 2003; 285:H1753-1758.
4. Baines CP, Song CX, Zheng YT, Wang GW, **Zhang J**, Wang OL, Guo Y, Bolli R, Cardwell EM, Ping P. Protein kinase C epsilon interacts with and inhibits the permeability transition pore in cardiac mitochondria. *Circ Res*. 2003; 92(8):873-880.
5. Edmondson RD, Vondriska TM, Biedermant KJ, **Zhang J**, Jones RC, Zheng YT, Allen DL, Xiu JX, Cardwell EM, Pisano MR, Ping P. Protein kinase C ϵ signaling complexes include metabolism-and transcription/translation-related proteins. *Mol Cell Proteomics*. 2002; 1:421-433.
6. Baines CP, **Zhang J**, Wang GW, Zheng YT, Xiu JX, Cardwell EM, Bolli R, Ping P. Mitochondrial PKC epsilon and MAPK form signaling modules in the murine heart: enhanced mitochondrial PKC epsilon-MAPK interactions and differential MAPK activation in PKC epsilon-induced cardioprotection. *Circ Res*. 2002; 90:390-397.
7. Ping P, Song C, **Zhang J**, Guo Y, Cao X, Li RC, Wu W, Vondriska TM, Pass JM, Tang XL, Pierce WM, Bolli R. Formation of protein kinase C(epsilon)-Lck signaling modules confers cardioprotection. *J Clin Invest*. 2002; 109:499-507.
8. Balafanova Z, Bolli R, **Zhang J**, Zheng Y, Pass JM, Bhatnagar A, Tang XL, Wang O, Cardwell E, Ping P. Nitric oxide induces nitration of PKC epsilon, facilitating PKC epsilon translocation via enhanced PKC epsilon-RACK2 interactions: A novel mechanism of NO-triggered activation of PKC epsilon. *J Biol Chem*. 2002; 277:15021-15027.
9. Song C, Vondriska TM, Wang GW, Klein JB, Cao X, **Zhang J**, Kang YJ, D'Souza S, Ping P. Molecular conformation dictates signaling module formation: example of PKC epsilon and Src tyrosine kinase. *Am J Physiol*. 2002; 282:H1166-1171.

10. Pass JM, Gao J, Jones WK, Wead WB, Wu X, **Zhang J**, Baines CP, Bolli R, Zheng YT, Joshua IG, Ping P. Enhanced PKC beta II translocation and PKC beta II-RACK1 interactions in PKC epsilon-induced heart failure: a role for RACK1 *Am J Physiol*. 2001; 281:H2500-2510.
11. **Zhang J**, Vondriska TM, Song C, Tang XL, Cao X, Baines CP, Pass JM, Wang S, Bolli R, Ping P. Protein kinase C epsilon-Src modules direct signal transduction in nitric oxide-induced cardioprotection: complex formation as a means for cardioprotective signaling. *Circ Res*. 2001; 88:1306-1313.
12. Pass JM, Zheng Y, Wead WB, **Zhang J**, Li RC, Bolli R, Ping P. PKC epsilon activation induces dichotomous cardiac phenotypes and modulates PKC epsilon-RACK interactions and RACK expression. *Am J Physiol*. 2001; 280:H946-955.
13. Ping P, **Zhang J**, Pierce WM Jr, Bolli R. Functional proteomic analysis of protein kinase C epsilon signaling complexes in the normal heart and during cardioprotection. *Circ Res*. 2001; 88:59-62.
14. Li RCX, Ping P, **Zhang J**, Wead WB, Cao X, Gao J, Zheng Y, Huang S, Han J, Bolli R. PKC epsilon modulates NF-kappa B and AP-1 via mitogen-activated protein kinases in adult rabbit cardiomyocytes. *Am J Physiol*. 2000; 279:H1679-1689.
15. Ping P, **Zhang J**, Huang S, Cao X, Tang XL, Li RC, Zheng YT, Qiu Y, Clerk A, Sugden P, Han J, Bolli R. PKC-dependent activation of p46/p54 JNKs during ischemic preconditioning in conscious rabbits. *Am J Physiol*. 1999; 277:H1771-1785.
16. Ping P, **Zhang J**, Zheng YT, Li RCX, Dawn B, Tang XL, Takano H, Balafanova Z, Bolli R. Demonstration of selective protein kinase C-dependent activation of Src and Lck tyrosine kinases during ischemic preconditioning in conscious rabbits. *Circ Res*. 1999; 85:542-550.
17. Ping P, **Zhang J**, Cao X, Li RCX, Kong D, Tang XL, Qiu Y, Manchikalapudi S, Auchampach JA, Black RG, Bolli R. PKC-dependent activation of p44/p42 MAPKs during myocardial ischemia-reperfusion in conscious rabbits. *Am J Physiol*. 1999; 276:H1468-1481.
18. Ping P, Takano H, **Zhang J**, Tang XL, Qiu Y, Li RCX, Banerjee S, Dawn B, Balafanova Z, Bolli R. Isoform-selective activation of protein kinase C by nitric oxide in the heart of conscious rabbits: a signaling mechanism for both nitric oxide-induced and ischemia-induced preconditioning. *Circ Res*. 1999; 84:587-604.
19. Bolli R, Dawn B, Tang XL, Qiu Y, Ping P, Xuan YT, Jones WK, Takano H, Guo Y, **Zhang J**. The nitric oxide hypothesis of late preconditioning. *Basic Res Cardiol*. 1998; 93:325-338.
20. Qiu Y, Ping P, Tang XL, Manchikalapudi S, Rizvi A, **Zhang J**, Takano H, Wu WJ, Teschner S, Bolli R. Direct evidence that protein kinase C plays an essential role in the development of late preconditioning against myocardial stunning in conscious rabbits and that epsilon is the isoform involved. *J Clin Invest*. 1998; 101:2182-2198.
21. Ping P, **Zhang J**, Qiu Y, Tang XL, Manchikalapudi S, Cao X, Bolli R. Ischemic preconditioning induces selective translocation of protein kinase C isoforms epsilon and eta in the heart of conscious rabbits without subcellular redistribution of total protein kinase C activity. *Circ Res*. 1997; 81:404-414.
22. **Zhang J**, Xu JL. Oleyl oleate synthesis by immobilized lipase from *Candida sp.*-1619. *Chinese J of Biotechnology*. 1995; 11:243-252.

23. Xu JL, **Zhang J**. Studies on the lipase from *Penicillium Camembertii*-PG3. *Acta Mycologica Sinica*. 1995; 14:136-142.
24. **Zhang J**, Xu JL. Oleyl oleate synthesis by immobilized lipase from *Candida sp.*-1619. *Chinese Journal of Biotechnology*. 1995; 11:325-331.

SELECTED ABSTRACTS

1. **Zhang J**, Zong C, Vondriska TM, Qiao X, Cardwell EM, Ping P. Proteomic mapping of cardiac protective signaling complexes using blue native electrophoresis and bioinformatic modeling. *FASEB J*. 2003; 17:A76.
2. Vondriska TM, Zhang S, **Zhang J**, Qiao X, Zheng YT, Cardwell EM, Ping P. Oracle regulates cardiac PKC ϵ signaling complexes in preconditioning. *FASEB J*. 2003; 17:A106.
3. Zhang S, Baines CP, **Zhang J**, Vondriska TM, Qiao X, Guo Y, Bhatnagar A, Bolli R, Ping P, Wang GW. Acrolein exacerbates ischemic injury in the heart and disrupts PKC ϵ protective signal transduction. *FASEB J*. 2003; 17:A880.
4. **Zhang J**, Vondriska TM, Wang GW, Baines CP, Stidam JM, Cardwell EM, Ping P. Characterization of a signaling complex by its protein composition, molecular architecture, and spatial distribution: defining the role of PKC ϵ -Akt-eNOS modules in cardioprotection via functional proteomics. *Circulation*. 2002; 106:II-6.
5. Ping P, **Zhang J**, Vondriska TM, Guo Y, Stidam JM, Cardwell EM, Wang GW, Bolli R. Cardiac-specific transgenesis of iNOS confers an infarct-sparing effect: signaling mechanisms involve site-specific iNOS tyrosine phosphorylation by Src tyrosine kinase. *Circulation*. 2002; 106: II-56.
6. Zheng YT, Vondriska TM, **Zhang J**, Edmondson RD, Ping P. Oracle, a PDZ and LIM domain-containing protein, particulate in PKC ϵ protective signaling complexes at distinct subcellular locations in the heart. *Circulation*. 2002; II-312.
7. Baines CP, Song C-X, Edmondson RD, Beiderman KJ, Jones RC, **Zhang J**, Cao X, Ping P. Functional proteomic analysis reveals an interaction between protein kinase C ϵ and the voltage-dependent anion channel in cardiac mitochondria. *FASEB J*. 2002; 16:A883.
8. **Zhang J**, Wang G-W, Baines CP, Zheng YT, Ping P. Protein kinase C ϵ forms a signaling module with Akt and eNOS: differential modulation by myocardial protection and myocardial toxicity. *FASEB J*. 2002; 16:A177.
9. **Zhang J**, Bolli R, Zheng YT, Xiu J, Wang GW, Pierce WM, Ping P. Subcellular location is a major determinant of PKC ϵ complex assembly: a view gained via functional proteomic analyses in cardioprotection. *Circulation*. 2001, 104: II-209.
10. Vondriska TM, Tang XL, **Zhang J**, Bolli R, Ping P. Bmx, a member of the Tec family of non-receptor tyrosine kinases, is a novel participant in the signaling system of NO-induced cardioprotection. *Circulation*. 2001, 104: II-61.
11. Baines CP, Wang GW, **Zhang J**, Zheng YT, Klein JB, Kang YJ, Bolli R, Ping P. Mitochondrial co-localization of MAPK with PKC ϵ in the murine heart: increased mitochondrial PKC ϵ -MAPK signaling module formation in PKC ϵ transgenic mice. *Circulation*. 2001, 104: II-102.

12. Vondriska TM, Ping P, **Zhang J**, Tang XL, Bolli R. PKC ϵ -dependent upregulation of PKC ϵ activity during nitric oxide-induced cardioprotection. **Circulation**. 2001, 104: II-208.
13. Song C, **Zhang J**, Cao X, Vondriska TM, Bolli R, Ping P. Conformation of molecules dictates signaling module formation: example of PKC ϵ and Src tyrosine kinase. **Circulation**. 2001, 104: II-210.
14. Baines CP, **Zhang J**, Zheng YT, Xiu JX, Bolli R, Ping P. Transgenic activation of PKC ϵ preferentially increases mitochondrial PKC ϵ expression and enhances formation of a mitochondrial PKC ϵ -JNK signaling complex in the mouse heart. **FASEB J**. 2001; 15:A103.
15. Vondriska TM, **Zhang J**, Tang XL, Bolli R, Ping P. Src tyrosine kinase forms a signaling complex with PKC ϵ during NO-induced preconditioning that is subcellular compartment-specific and PKC ϵ activation-dependent. **FASEB J**. 2001; 15:A119.
16. Vondriska TM, Edmondson R, **Zhang J**, Biederman K, Zheng Y, Jones R, and Ping P. Functional proteomics: PKC ϵ -mediated cardioprotective signaling involves metabolism- and transcription/translation-related proteins. *AHA Myocardial Ischemia Meeting*. 2001.
17. Vondriska TM, Song C, Wang G, Cao X, Zhang J, D'Souza S, Bolli R, and Ping P. Molecular conformation determines protein-protein interactions between PKC ϵ and Src tyrosine kinase. 2001. *AHA Myocardial Ischemia Meeting*. 2001.
18. Wang GW, **Zhang J**, Baines CP, Song CX, Zheng YT, Klein JB, Kang YJ, Bolli R, Ping P. Cardioprotection involves subcellular compartment-dependent localization of PKC ϵ -Src and PKC ϵ -Lck signaling modules. **J Mol Cell Cardiol**. 2001.
19. Vondriska TM, **Zhang J**, Song C, Tang XL, Bolli R, Ping P., PKC ϵ -Src modules direct signal transduction in NO-induced cardioprotection: Complex formation as a means for cardioprotection. **J Mol Cell Cardiol** (suppl.), 2001
20. Song CX, Cao X, **Zhang J**, Vondriska TM, Bolli R, Ping P. PKC ϵ binds to and activated Src tyrosine kinase *in vitro*: Characterizing a cardiac PKC ϵ -Src module. **J Mol Cell Cardiol** (suppl.), 2001.
21. Balafanova Z, Gao JM, **Zhang J**, Bolli R, Ping P. Nitric oxide induces activation of PKC ϵ via a cGMP-dependent mechanism in adult rabbit cardiac myocytes. **J Mol Cell Cardiol**. 2001; 33:A7.
22. Pass J, **Zhang J**, Zheng YT, Bolli R, Ping P. Enhanced PKC β II translocation and PKC β II-RACK1 interactions in PKC ϵ -induced cardiac hypertrophy and failure: a role for RACK1. **J Mol Cell Cardiol**. 2001; 33:A90.
23. Ping P, **Zhang J**, Zheng YT, Li CX, Guo YR, Bao WK, Bolli R. Cardiac targeted transgenesis of active PKC epsilon renders the heart resistant to infarction. **Circulation** 2000, 102: II-24.
24. Ping P, Guo YR, **Zhang J**, Li CX, Bao WK, Wu WJ, Bolli R. Targeted deletion of Lck tyrosine kinase abrogates the NF-kappa B activation and the infarct-sparing effect induced by late preconditioning. **Circulation** 2000, 102: II-211.
25. Song C, **Zhang J**, Gao JM, Zheng YT, Cao X, Wead WB, Bolli R, Ping P. Lck tyrosine kinase is a direct substrate of PKC epsilon in cardiac myocytes. **Circulation** 2000, 102: II-211.

26. Pass JM, Zheng YT, **Zhang J**, Li CX, Bolli R, Ping P. Dichotomous cardiac phenotypes are congruous with differential interactions between protein kinase C epsilon and its receptor proteins. **Circulation** 2000, 102: II-160.
27. Li CX, **Zhang J**, Zheng YT, Wead WB, Gao JM, Cao X, Bolli R, Ping P. PKC epsilon enhances NF-kappa B activation via modulation of IKK complex in cardiac myocytes. **Circulation** 2000, 102: II-120.
28. Ping P, **Zhang J**, Bolli R. Functional proteomic analysis of protein kinase C ϵ signaling complexes in preconditioning. **Circulation** 2000, 102: II-221.
29. Pass JM, **Zhang J**, Li CX, Zheng YT, Bolli R, Ping P. Stoichiometric analyses of PKC and RACK expression in the FVB/N mouse heart. **J Mol Cell Cardiol.** 2000, 32: A25.
30. Gao JM, Zheng YT, **Zhang J**, Bolli R, Li CX, Wead WB, Ping P. Transgenic overexpression of the η isoform of PKC does not trigger cardiac hypertrophy. **J Mol Cell Cardiol.** 2000; 32: A58.
31. **Zhang J**, Bolli R, Tang XL, Jones WK, Ping P. Direct demonstration of both tyrosine and serine phosphorylation of I κ B during ischemic preconditioning in conscious rabbits. **Circulation** 1999, 100: I-56.
32. **Zhang J**, Bolli R, Lalli J, Tang XL, Li CX, Zheng YT, Pass JM, Ping P. Ischemic preconditioning and phorbol ester redistribute protein kinase C ϵ to the nucleus, sarcolemmal membranes, and mitochondria in rabbit myocardium. **Circulation** 1999, 100: I-325.
33. Li CX, Ping P, **Zhang J**, Joshua IG, Zheng YT, Gao JM, Bolli R. Protein kinase C epsilon activates nuclear factor-KappaB through a Src-Dependent signaling pathway in adult rabbit cardiac myocytes. **Circulation** 1999, 100: I-325.
34. Li CX, Ping P, **Zhang J**, Cao X, Zheng YT, Gao JM, Xuan YT, Wead WB, Joshua IG, Bolli R. PKC ϵ activates NF κ B and AP-1 via a ERK-dependent signaling pathway in the adult rabbit cardiac myocytes. **J Mol Cell Cardiol.** 1999, 31: A18.
35. **Zhang J**, Wead WB, Jones WK, Wu X, Gao JM, Kong DY, Li CX, Zheng YT, Ping P. Activation of PKC ϵ induces hypertrophy and heart failure in a dose-dependent fashion in mice. **J Mol Cell Cardiol.** 1999, 31: A18.
36. Pass JM, **Zhang J**, Zheng YT, Li CX, Ping P. RACK expression is coordinated with PKC ϵ expression in the mouse heart. **J Mol Cell Cardiol.** 1999, 31: A19.
37. Ping P, Li RCX, **Zhang J**, Kong D, Jones WK, Zheng YT, Cao X, Bolli R. Identification of two distinct mechanisms for the activation of p44/p42 MAPKs in vivo by a PKC epsilon dependent signaling pathway in transgenic mice. **Circulation** 1998, 98:I-416.
38. Ping P, **Zhang J**, Li CX, Tang XL, Qiu Y, Banerjee S, Zheng YT, Bolli R. Ischemic preconditioning (PC) induces selective activation of Src and Lck tyrosine kinases in conscious rabbits via a PKC-dependent pathway. **Circulation** 1998, 98:I-71.
39. Ping P, **Zhang J**, Cao X, Kong D, Tang XL, Qiu Y, Manchikalapudi S, Li CX, Bolli R. Brief episodes of ischemia induce activation of the MEK1/2-p44/p42 MAPK signaling cascade in the cytosolic fraction and subsequent nuclear translocation of p44/p42 MAPKs via a PKC-dependent pathway in conscious rabbits. **Circulation** 1998; 98:I-586.

40. Ping P, Kong D, Jones WK, Li CX, **Zhang J**, Cao X, Wu X, Wead WB, Harris PD, Joshua IG, Bolli R. Cardiac targeted expression of PKC epsilon isoform selectively alters PKC activity in the mouse heart. **FASEB J**. 1998, 12:A-75.
41. Ping P, **Zhang J**, Cao X, Qiu Y, Tang XL, Clerk A, Sugden P, Bolli R. Activation of the P38MAPK and the p46/p54 JNKs after brief episodes of ischemia/reperfusion in the heart of conscious rabbits. **J Mol Cell Cardiol**. 1998, 30: A-263.
42. Ping P, Cao X, Kong D, **Zhang J**, Li CX, Yang Y, Bolli R. PKC epsilon isoform induces activation of the p42/p44 MAPKs and the p46/p54 JNKs in adult rabbit cardiac myocytes. **J Mol Cell Cardiol**. 1998; 30:A-263.
43. Ping P, **Zhang J**, Qiu Y, Tang XL, Manchikalapudi S, Bolli R. Repetitive episodes of myocardial ischemia and reperfusion induce translocation of protein kinase C epsilon isoform in conscious rabbits, which is associated with late preconditioning against myocardial stunning. **Circulation** 1996; 94:I-660.
44. Xu JL, **Zhang J**. Lipase catalyzed synthesis of 2-ethyl hexanol fatty acid esters. Oral presentation, the fourth China-Japan Joint symposium on Enzyme Engineering, Oct., 1996.
45. **Zhang J**, Jia-Li Xu. Hydrolysis and Synthesis of Oleyl Oleate Catalyzed by Lipases. Poster presentation, the Second National Conference on Enzyme Engineering, **Acta Scientiarum Naturalium Universitatis Jilinensis** 1993, P4-6.
46. DD Gu, J.I. Xu, **Zhang J**. Immobilization of *Candida sp.-1619* Lipase for Synthesis wax-esters. Poster presentation, the Second National Conference on Enzyme Engineering, **Acta Scientiarum Naturalium Universitatis Jilinensis** 1993, P4-7.

BOOK CHAPTERS:

1. Pass PM, **Zhang J**, Vondrisk TM, Ping P. (Newton AC eds.) Functional proteomic analysis of the protein kinase C signaling system. *Protein kinase C protocols*, Methods in molecular biology, 233: 369-385, Humana Press, 2003.
2. Vondrisk TM, **Zhang J**, Ping P. (Van Eyk JE and Dunn MJ, eds.) Proteomic characterization of protein kinase C signaling tasks in the hearts. *Proteomic and genomic analysis of cardiovascular disease*. Wiley-VCH, 2003.
3. Baines CP, **Zhang J**, Ping P. Characterization of mitochondrial signaling complexes. Ping P, Hanash S, eds. *Functional proteomics: principles and practical applications*. The Humana Press Inc. (in press).
4. Rokosh G and **Zhang J** Co-editors. *Cardiac genes expression and regulation*. Methods in molecular biology, Humana Press. (in preparation)

NATIONAL MEETING ORAL PRESENTATION

Oral presentation: American Heart Association 75th Scientific Sessions, Chicago, IL. Title: Characterization of a signaling complex by its protein composition, molecular architecture, and spatial distribution: defining the role of PKC ϵ -Akt-eNOS modules in cardioprotection via functional proteomics. November 11, 2002.